



# Simultaneous model-based clustering and visualization in the Fisher discriminative subspace

Charles Bouveyron, Camille Brunet

## ► To cite this version:

Charles Bouveyron, Camille Brunet. Simultaneous model-based clustering and visualization in the Fisher discriminative subspace. 2010. hal-00492406v1

**HAL Id: hal-00492406**

**<https://paris1.hal.science/hal-00492406v1>**

Preprint submitted on 15 Jun 2010 (v1), last revised 19 Apr 2011 (v4)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Simultaneous model-based clustering and visualization in the Fisher discriminative subspace

Charles BOUVEYRON<sup>1</sup> & Camille BRUNET<sup>2</sup>

<sup>1</sup> Laboratoire SAMM, EA 4543, Université Paris 1 Panthéon-Sorbonne  
90 rue de Tolbiac, 75013 Paris, France  
Email: `charles.bouveyron@univ-paris1.fr`

<sup>2</sup> IBISC, TADIB, FRE CNRS 3190, Université d'Evry Val d'Essonne  
40 rue de Pelvoux, CE 1455, 91020 Evry Courcouronnes, France  
Email: `camille.brunet@ibisc.univ-evry.fr`

## Abstract

Clustering in high-dimensional spaces is nowadays a recurrent problem in many scientific domains but remains a difficult task from both the clustering accuracy and the result understanding points of view. This paper presents a discriminative latent mixture (DLM) model which models the data in a latent orthonormal discriminative subspace with an intrinsic dimension lower than the dimension of the original space. By constraining model parameters within and between groups, a family of 8 parsimonious DLM models is exhibited and this allows to fit onto various situations. An estimation algorithm, called the Fisher-EM algorithm, is also proposed for estimating both the mixture parameters and the discriminative subspace. Experiments on simulated and real datasets show that the proposed approach outperforms existing clustering methods and provides a useful representation of the clustered data. The method is as well applied to the clustering of mass spectrometry data.

**Keywords:** high-dimensional clustering, model-based clustering, discriminative subspace, Fisher criterion, visualization, parsimonious models.

## 1 Introduction

In many scientific domains, the measured observations are nowadays high-dimensional and clustering such data remains a challenging problem. Indeed, the most popular clustering methods, which are based on the mixture model, show a disappointing behavior in high-dimensional spaces. They suffer from the well-known *curse of dimensionality* [6] which is mainly due to the

fact that model-based clustering methods are over-parametrized in high-dimensional spaces. Furthermore, in several applications such as mass spectrometry or genomics, the number of available observations is small compared to the number of variables and such a situation increases the difficulty of the problem.

Hopefully, since the dimension of observed data is usually higher than their intrinsic dimension, it is theoretically possible to reduce the dimension of the original space without loosing any information. Therefore, dimension reduction methods are traditionally used before the clustering step. Feature extraction methods such as Principal Component Analysis (PCA) or feature selection methods are very popular. However, these approaches of dimension reduction do not consider the classification task and provide a sub-optimal data representation for the clustering step. Indeed, dimension reduction methods imply an information loss which could be discriminative. Only few approaches combine dimension reduction with the classification aim but, unfortunately, those approaches are all supervised methods. Fisher Discriminant Analysis (FDA) (see Chap. 4 of [25]) is one of them in the supervised classification framework. FDA is a powerful tool for finding the subspace which best discriminates the classes and reveals the structure of the data. This subspace is spanned by the discriminative axes which maximize the ratio of the between class variance and the within class variance.

To avoid dimension reduction, several subspace clustering methods have been proposed in the past few years to model the data of each group in low-dimensional subspaces. These methods turned out to be very efficient in practice. However, since these methods model each group in a specific subspace, they are not able to provide a global visualization of the clustered data which could be helpful for the practitioner. Indeed, the clustering results of high-dimensional data are difficult to understand without a visualization of the clustered data. In addition, in scientific fields such as genomics or economics, original variables have an actual meaning and the practitioner could be interested in interpreting the clustering results according to the variable meaning.

In order to both overcome the curse of dimensionality and improve the understanding of the clustering results, this work proposes a method which adapts the traditional mixture model for modeling and classifying data in a discriminative subspace. For this, the proposed discriminative latent mixture (DLM) model combines the model-based clustering goals with the discriminative criterion introduced by Fisher. The estimation procedure proposed in this paper and named Fisher-EM has three main objectives: firstly, it aims to improve clustering performances with the use of a discriminative subspace, secondly, it avoids estimation problems linked to high dimensions through model parsimony and, finally, it provides a low-dimensional discriminative representation of the clustered data.

The reminder of this manuscript has the following organization. Section 2 reviews the problem of high-dimensional data clustering and existing solutions. Section 3 introduces the discriminative latent mixture model and its submodels. The link with existing approaches is also discussed in Section 3. Section 4 presents an EM-based procedure, called Fisher-

EM, for estimating the parameters of the DLM model. Initialization, model selection and computational issues are also considered in Section 4. In Section 5, the Fisher-EM algorithm is compared to existing clustering methods on simulated and real datasets. Section 6 presents the application of the Fisher-EM algorithm to a real-world clustering problem in mass-spectrometry imaging. Some concluding remarks and ideas for further works are finally given in Section 7.

## 2 Related works

Clustering is a traditional statistical problem which aims to divide a set of observations  $\{y_1, \dots, y_n\}$  described by  $p$  variables into  $K$  homogeneous groups. The problem of clustering has been widely studied for years and the reader could refer to [17, 27] for reviews on the clustering problem. However, the interest in clustering is still increasing since more and more scientific fields require to cluster high-dimensional data. Moreover, such a task remains very difficult since clustering methods suffer from the well-known *curse of dimensionality* [6]. Conversely, the *empty space phenomenon* [45], which refers to the fact that high-dimensional data do not fit the whole observation space but live in low-dimensional subspaces, gives hope to efficiently classify high-dimensional data. This section firstly reviews the framework of model-based clustering before exposing the existing approaches to deal with the problem of high dimension in clustering.

### 2.1 Model-based clustering and high-dimensional data

Model-based clustering, which has been widely studied by [17, 36] in particular, aims to partition observed data into several groups which are modeled separately. The overall population is considered as a mixture of these groups. Let us consider a given dataset of  $n$  observations  $\{y_1, \dots, y_n\}$  to divide into  $K$  homogeneous groups. Let us also assume that the observations  $\{y_1, \dots, y_n\}$  are independent realizations of a random vector  $Y \in \mathbb{R}^p$  and that  $f$  is its density function. Then, the finite mixture model is:

$$f(y) = \sum_{k=1}^K \pi_k f_k(y),$$

where  $f_k$  and  $\pi_k$  respectively represent the conditional density function and the mixture proportion of the  $k$ th component. The clusters are often modeled by the same parametric density function in which case the finite mixture model is:

$$f(y) = \sum_{k=1}^K \pi_k f(y; \theta_k),$$

where  $\theta_k$  is the parameter vector for the  $k$ th component. Moreover, the density function  $f_k$  is, most commonly, a multivariate Gaussian density  $\phi_k$  parametrized by its mean  $\mu_k$  and its covariance matrix  $\Sigma_k$ , such that the density function of  $Y$  can be written in this way

$$f(y) = \sum_{k=1}^K \pi_k \phi(y; \mu_k, \Sigma_k).$$

The unknown parameters  $\pi_k$ ,  $\mu_k$  and  $\Sigma_k$ ,  $k = 1, \dots, K$ , are usually estimated by the maximum likelihood owing to the Expectation-Maximization (EM) algorithm proposed by [12] and extended by [10], [11], [35]. Since the maximization of the complete likelihood is intractable, the EM algorithm is based on the maximization of the conditional expectation of the likelihood given current parameters and consists of two steps: the expectation step first computes the posterior probabilities and the second step maximizes the expectation of the complete likelihood subject to the posterior probabilities. These two steps are iteratively computed until the convergence of the likelihood. The partition is designed afterward using the maximum a posteriori (MAP) rule: each observation is assigned to the group with the highest posterior probability.

Unfortunately, model-based clustering methods show a disappointing behavior when the number of observations is small compared to the number of parameters to estimate. Indeed, in the case of the Gaussian mixture model, the number of parameters to estimate is a function of the square of the dimension  $p$  and the estimation of this potentially large number of parameters is consequently difficult with a small dataset. In particular, when the number of observations  $n$  is of the same order than the number of dimensions  $p$ , model-based clustering methods have to face numerical problems due to the ill-conditioning of the covariance matrices. Furthermore, it is not possible to use the Gaussian mixture model without restrictive assumptions for clustering a dataset for which  $n$  is smaller than  $p$ . Indeed, for clustering such data, it would be necessary to invert  $K$  covariance matrices which would be singular. To overcome these problems, several strategies have been proposed in the literature among which dimension reduction and subspace clustering.

## 2.2 Dimension reduction and clustering

Earliest approaches proposed to overcome the problem of high dimension in clustering by reducing the dimension before using a traditional clustering method. Among the unsupervised tools of dimension reduction, Principal Component Analysis (PCA) [29] is the traditional and certainly the most used technique for dimension reduction. It aims to project the data on a lower dimensional subspace in which axes are build by maximizing the variance of the projected data. However, PCA is a linear tool, which means that non-linear dependencies are not take into account. Therefore, other alternatives have been proposed: Kambhatla and Leen [30] for example, have developed a method which uses PCA but locally, in restricted

parts of the space; Scholkopf and Smola [43] have proposed a method called Kernel PCA which transforms the original data in a higher dimensional space before applying PCA in these transformed data; Hastie *et al.* [24] and Girard [20] have also proposed non-linear versions of PCA. Other non-linear projection tools can be mentioned such as Kohonen’s maps [31]. For a review of these dimension reduction approaches, see [48]. An other way to reduce the dimension in an unsupervised problem has been recently considered in [41] and [34] in which the problem of feature selection for model-based clustering is recasted as a model selection problem. However, such approaches remove variables and consequently information which could have been discriminative for the clustering task.

### 2.3 Subspace clustering

In the past few years, new approaches focused on the modeling of each group in specific subspaces of low dimensionality. Subspace clustering methods can be split into two categories: heuristic and probabilistic methods. Heuristic methods use algorithms to search for subspaces of high density within the original space. On the one hand, bottom-up algorithms use histograms for selecting the variables which best discriminate the groups. The Clique algorithm [1] was one of the first bottom-up algorithms and remains a reference in this family of methods. On the other hand, top-down algorithms use iterative techniques which start with all original variables and remove at each iteration the dimensions without groups. A review on heuristic methods is available in [40]. Conversely, probabilistic methods assume that the data of each group live in a low-dimensional latent space and usually model the data with a generative model. Earlier strategies [42] are based on the factor analysis model which assumes that the latent space is related with the observation space through a linear relationship. This model was recently extended in [5, 37] and yields in particular the well known mixture of probabilistic principal component analyzers [46]. Recent works [9, 38] proposed two families of parsimonious and regularized Gaussian models which partially encompass previous approaches. All these techniques turned out to be very efficient in practice to cluster high-dimensional data. However, despite their qualities, these probabilistic methods mainly consider the clustering aim and do not take enough into account the visualization and understanding aspects.

### 2.4 From Fisher’s theory to discriminative clustering

In the case of supervised classification, Fisher poses, in his precursor work [15], the problem of the discrimination of three species of iris described by four measurements in these terms: *What linear function of the four measurements will maximize the ratio of the difference between the specific means to the standard deviation within species?* Such a remark has led to useful approaches in supervised classification and in dimension reduction.

### 2.4.1 Fisher discriminant analysis

The main goal of Fisher was to find a linear subspace that separates the classes according to a criterion (see [14] for more details). For this, Fisher assumes that the dimensionality  $p$  of the original space is greater than the number  $K$  of classes. Fisher discriminant subspace looks for a linear transformation  $U$  which projects the observations in a discriminant and low dimensional subspace of dimension  $d$  such that the linear transformation  $U$  of dimension  $p \times d$  aims to maximize a criterion which is large when the between-class covariance matrix ( $S_B$ ) is large and when the within-covariance matrix ( $S_W$ ) is small. Four different criteria can be found in the literature which verify such a constraint (see [19] for a review). The criterion which is traditionally used is:

$$J(U) = \text{tr} \left( (U^t S_W U)^{-1} U^t S_B U \right),$$

where  $\text{tr}(\cdot)$  is the trace operator and:

$$S_W = \frac{1}{n} \sum_{k=1}^K \sum_{i \in C_k} (y_i - m_k)^T (y_i - m_k)$$

$$S_B = \frac{1}{n} \sum_{k=1}^K n_k (m_k - \bar{y})^T (m_k - \bar{y})$$

are respectively the within and the between covariance matrices,  $m_k = \frac{1}{n_k} \sum_{i \in C_k}^K y_i$  is the empirical mean of the observations  $y_i$  in the class  $k$  and  $\bar{y} = \frac{1}{n} \sum_{k=1}^K n_k m_k$  is the mean of the observations. This criterion has to be maximized to reach the solution expected by Fisher and the solution of this optimization problem is the eigenvectors associated to the  $K - 1$  largest eigenvalues of the matrix  $S_W^{-1} S_B$ . Once the discriminative axes determined, linear discriminant analysis (LDA) is usually applied to classify the data into this subspace.

### 2.4.2 Regularization of Fisher discriminant analysis

The optimization of the Fisher criterion supposes the non singularity of the matrix  $S_W$  but it appears that the singularity of  $S_W$  occurs frequently, particularly in the case of very high dimensional space or in the case of under-sampled problems. In the literature, different solutions are proposed to deal with such a problem in a supervised classification framework. The regularized discriminant analysis (RDA), proposed by Friedman [18], suggests that the covariance matrix depends on two regularized parameters ; the pseudo-inverse LDA [19] proposes to use the *pseudo-inverse* of the within covariance matrix and also the penalized discriminant analysis (PDA) [23] which deals with high correlated variables or under-sampled cases. More recently, Howland [26] used the generalized singular value decomposition (GSVD) to overcome the singularity problem of the pooled scatter matrix ( $S_W$ ) in using a special shape of the within and between covariance matrices and Zhang [52] improves this approach es-

pecially in the case of small observations but large set of variables. An other approach has been suggested by Jin *et al.* [28], namely the uncorrelated LDA (ULDA) approach, to deal with under-sampled problems and has the property that the features in the reduced space are uncorrelated.

#### 2.4.3 Extension to unsupervised classification: discriminative clustering

Since clustering approaches are sensitive to high-dimensional and noisy data, recent works focused on combining low dimensional discriminative subspace with one of the most used clustering algorithm: kmeans. The proposed methods iteratively compute a discriminative subspace based on the Fisher criterion given the previous partition and obtains a new partition by kmeans subject to this subspace. The first basic algorithm has been proposed by Xu *et al.* [49] and has then been extended by De la Torre [32] who develops a discriminative cluster analysis (DCA) method in the case of non invertible scatter matrix. A theoretical framework is suggested by Ding in [13] when both tasks perform simultaneously since Fisher discriminant analysis and kmeans clustering optimize the same objective function. More recently, Ye *et al.* [51] reformulate the iterative problem of clustering in discriminative subspace and they show that the iterative subspace selection and kmeans clustering is equivalent to kernel kmeans task with a specific kernel Gram matrix. However, these approaches do not really compute the discriminant subspace and are not interested in the visualization and the understanding of the data.

### 3 Model-based clustering in a discriminative subspace

This section introduces a mixture model, called the discriminative latent mixture model, which ambitions to find both a parsimonious and discriminative fit for the data in order to ease the clustering and the visualization of the clustered data. The modeling proposed in this section is mainly based on two key-ideas: firstly, actual data are assumed to live in a latent subspace with an intrinsic dimension lower than the dimension of the observed data and, secondly, a subspace of  $K - 1$  dimensions is theoretically sufficient to discriminate  $K$  groups.

#### 3.1 The discriminative latent mixture model

Let  $\{y_1, \dots, y_n\} \in \mathbb{R}^p$  denote a dataset of  $n$  observations that one wants to cluster into  $K$  homogeneous groups, *i.e.* adjoin to each observation  $y_j$  a value  $z_j \in \{1, \dots, K\}$  where  $z_i = k$  indicates that the observation  $y_i$  belongs to the  $k^{th}$  group. On the one hand, let us assume that  $\{y_1, \dots, y_n\}$  are independent observed realizations of a random vector  $Y \in \mathbb{R}^p$  and that  $\{z_1, \dots, z_n\}$  are also independent realizations of a random vector  $Z \in \{1, \dots, K\}$ . On the other hand, let  $\mathbb{E} \subset \mathbb{R}^p$  denote a latent space assumed to be the most discriminative subspace of dimension  $d = K - 1$  such that  $\mathbf{0} \in \mathbb{E}$  and where  $d$  is strictly lower than the dimension  $p$  of



the observed space. Moreover, let  $\{x_1, \dots, x_n\} \in \mathbb{E}$  denote the actual data, described in the latent space  $\mathbb{E}$  of dimension  $d$ , which are in addition presumed to be independent unobserved realizations of a random vector  $X \in \mathbb{E}$ . Finally, for each group, the observed variable  $Y \in \mathbb{R}^p$  and the latent variable  $X \in \mathbb{E}$  are assumed to be linked through a linear transformation:

$$Y|_{Z=k} = UX + \varepsilon, \quad (3.1)$$

where  $d < p$ ,  $U$  is the  $p \times d$  orthogonal matrix such as  $U^T U = I_d$  and  $\varepsilon \in \mathbb{R}^p$  is a centered Gaussian noise term with covariance matrix  $\Psi_k$ , for  $k = 1, \dots, K$ :

$$\varepsilon|Z = k \sim \mathcal{N}(\mathbf{0}, \Psi_k).$$

Following the classical framework of model-based clustering, each group is in addition assumed to be distributed according to a Gaussian density function within the latent space  $\mathbb{E}$ . Hence, the random vector  $X \in \mathbb{E}$  has the following conditional density function:

$$X|Z = k \sim \mathcal{N}(\mu_k, \Sigma_k),$$

where  $\mu_k \in \mathbb{R}^d$  and  $\Sigma_k \in \mathbb{R}^{d \times d}$  are respectively the mean and the covariance matrix of the  $k$ th group. Since the latent space  $\mathbb{E}$  is assumed to have an orthonormal basis,  $\Sigma_k$  is a diagonal matrix and can be written  $\Sigma_k = \text{diag}(\alpha_{k1}, \dots, \alpha_{kd})$ . With these assumptions and conditionally to  $X$  and  $Z$ , the random vector  $Y \in \mathbb{R}^p$  has the following conditional distribution:

$$Y|X, Z = k \sim \mathcal{N}(U\mu_k, \Psi_k),$$

and its marginal distribution is therefore a mixture of Gaussians:

$$f(y) = \sum_{k=1}^K \pi_k \phi(y; m_k, S_k),$$

where  $\pi_k$  is the mixture proportion of the  $k$ th group and:

$$\begin{aligned} m_k &= U\mu_k, \\ S_k &= U\Sigma_k U^T + \Psi_k, \end{aligned}$$

are respectively the mean and the covariance matrix of the  $k$ th group in the observation space. Let us also define  $W = [U, V]$  a  $p \times p$  matrix which verifies  $W^T W = W W^T = I_p$  and for which  $V$  is the orthonormal complement of  $U$  defined above. We finally assume that the noise




Figure 1: Graphical summary of the  $\text{DLM}_{[\alpha_{kj}\beta_k]}$  model

covariance matrix  $\Psi_k$  is such that  $\Delta_k = W^T S_k W$  has the following form:

$$\Delta_k = \left( \begin{array}{cc|cc} \alpha_{k1} & 0 & & \\ & \ddots & & \\ 0 & \alpha_{kd} & & \\ \hline & & \beta_k & 0 \\ & & \ddots & \ddots \\ \mathbf{0} & & 0 & \beta_k \end{array} \right) \left\{ \begin{array}{l} d = K - 1 \\ (p - d) \end{array} \right.$$

This model, called the discriminative latent mixture (DLM) model and referred to by  $\text{DLM}_{[\alpha_{kj}\beta_k]}$  in the sequel, is summarized by Figure 1. The  $\text{DLM}_{[\alpha_{kj}\beta_k]}$  model is therefore parametrized by the parameters  $\pi_k$ ,  $\mu_k$ ,  $U$ ,  $\alpha_{kj}$  and  $\beta_k$ , for  $k = 1, \dots, K$  and  $j = 1, \dots, d$ . On the one hand, the mixture proportions  $\pi_1, \dots, \pi_K$  and the means  $\mu_1, \dots, \mu_K$  parametrize in a classical way the prior probability and the average latent position of each group respectively. On the other hand,  $U$  defines the latent subspace  $\mathbb{E}$  by parametrizing its orientation according to the basis of the original space. Finally,  $\alpha_{k1}, \dots, \alpha_{kd}$  parametrize the variance of the  $k$ th group within the latent subspace  $\mathbb{E}$  whereas  $\beta_k$  parametrizes the variance of this group outside  $\mathbb{E}$ . With these notations and from a practical point of view, the variance of the actual data is therefore modeled by  $\alpha_{k1}, \dots, \alpha_{kd}$  and the variance of the noise is modeled by  $\beta_k$ .

### 3.2 The submodels of the $\text{DLM}_{[\alpha_{kj}\beta_k]}$ model

Starting with the  $\text{DLM}_{[\alpha_{kj}\beta_k]}$  model presented in the previous paragraph, several submodels can be generated by applying constraints on parameters of the diagonal matrix  $\Delta_k$ . For in-

Model	Nb. of parameters	$K = 4$ and $p = 100$
$\text{DLM}_{[\alpha_{kj}\beta_k]}$	$(K - 1) + Kp + (K - 1)(p - K/2) + K^2$	713
$\text{DLM}_{[\alpha_{kj}\beta]}$	$(K - 1) + Kp + (K - 1)(p - K/2) + K(K - 1) + 1$	710
$\text{DLM}_{[\alpha_k\beta_k]}$	$(K - 1) + Kp + (K - 1)(p - K/2) + 2K$	705
$\text{DLM}_{[\alpha_k\beta]}$	$(K - 1) + Kp + (K - 1)(p - K/2) + K + 1$	702
$\text{DLM}_{[\alpha_j\beta_k]}$	$(K - 1) + Kp + (K - 1)(p - K/2) + (K - 1) + K$	704
$\text{DLM}_{[\alpha_j\beta]}$	$(K - 1) + Kp + (K - 1)(p - K/2) + (K - 1) + 1$	701
$\text{DLM}_{[\alpha\beta_k]}$	$(K - 1) + Kp + (K - 1)(p - K/2) + K + 1$	702
$\text{DLM}_{[\alpha\beta]}$	$(K - 1) + Kp + (K - 1)(p - K/2) + 2$	698
Full-GMM	$(K - 1) + Kp + Kp(p + 1)/2$	20603
Com-GMM	$(K - 1) + Kp + p(p + 1)/2$	5453
Mixt-PPCA	$(K - 1) + Kp + K(d(p - (d + 1)/2) + d + 1) + 1$	4228 ( $d = 10$ )
Diag-GMM	$(K - 1) + Kp + Kp$	803
Sphe-GMM	$(K - 1) + Kp + K$	407

Table 1: Number of free parameters to estimate for the DLM models and some classical models (see text for details).

stance, the  $p - d$  last values of  $\Delta_k$  can be assumed to be common for the  $k$  classes, *i.e.*  $\beta_k = \beta$ ,  $\forall k = 1, \dots, K$ , meaning that the variance outside the discriminant subspace is common to all groups. This assumption can be viewed as modeling the noise variance with a unique parameter which seems natural for data obtained in a common acquisition process. Following the notation system introduces above, this submodel will be referred to by  $\text{DLM}_{[\alpha_{kj}\beta]}$ . The variance within the latent subspace  $\mathbb{E}$  can also be assumed to be isotropic for each group and the associated submodel is  $\text{DLM}_{[\alpha_k\beta_k]}$ . In this case, the variance of the data is assumed to be isotropic both within  $\mathbb{E}$  and outside  $\mathbb{E}$ . Similarly, it is possible to constrain the previous model to have the parameters  $\beta_k$  common between classes and this gives rise to the model  $\text{DLM}_{[\alpha_k\beta]}$ . Finally, the variance within the subspace  $\mathbb{E}$  can be assumed to be independent from the mixture component and this corresponds to the models  $\text{DLM}_{[\alpha_j\beta_k]}$ ,  $\text{DLM}_{[\alpha_j\beta]}$ ,  $\text{DLM}_{[\alpha\beta_k]}$  and  $\text{DLM}_{[\alpha\beta]}$ . We therefore enumerate 8 different DLM models and an overview of them is proposed in Table 1. The table also gives the number of free parameters to estimate according to  $K$  and  $p$  for the 8 DLM models and for some classical models. The Full-GMM model refers to the classical Gaussian mixture model with full covariance matrices, the Com-GMM model refers to the Gaussian mixture model for which the covariance matrices are assumed to be equal to a common covariance matrix ( $S_k = S$ ,  $\forall k$ ), Diag-GMM refers to

the Gaussian mixture model for which  $S_k = \text{diag}(s_{k1}^2, \dots, s_{kp}^2)$  with  $s_k^2 \in \mathbb{R}^p$  and Sphe-GMM refers to the Gaussian mixture model for which  $S_k = s_k^2 I_p$  with  $s_k^2 \in \mathbb{R}$ . Finally, Mixt-PPCA denotes the subspace clustering model proposed by Tipping and Bishop in [46]. In addition to the number of free parameters to estimate, Table 1 gives this number for specific values of  $K$  and  $p$  in the right column. The number of free parameters to estimate given in the central column can be decomposed in the number of parameters to estimate for the proportions ( $K - 1$ ), for the means ( $Kp$ ) and for the covariance matrices (last terms). Among the classical models, the Full-GMM model is a highly parametrized model and requires the estimation of 20603 parameters when  $K = 4$  and  $p = 100$ . Conversely, the Diag-GMM and Sphe-GMM model are very parsimonious models since they respectively require the estimation of only 803 and 407 parameters when  $K = 4$  and  $p = 100$ . The Com-GMM and Mixt-PPCA models appear to both have an intermediate complexity. However, the Mixt-PPCA model is a less constrained model compared to the Diag-GMM model and should be preferred for clustering high-dimensional data. Finally, the DLM models turn out to have a low complexity whereas their modeling capacity is comparable to the one of the Mixt-PPCA model. In addition, the complexity of the DLM models depends only from  $K$  and  $p$  whereas the Mixt-PPCA model depends from an hyper-parameter  $d$ .

### 3.3 Links with existing models

At this point, some links can be established with models existing in the literature. The closest models have been proposed in [5], [9] and [38]. First, in [9], the authors proposed a family of 28 parsimonious and flexible Gaussian models ranging from a very general model, referred to as  $[a_{kj}b_kQ_kd_k]$ , to very simple models. Among this family of parsimonious models, 14 models assume that the orientation of the group-specific subspaces is common (common  $Q_k$ ). In particular, this work can be viewed as an extension of the mixture of principal component analyzer (Mixt-PPCA) model [46]. The following year, McNicholas and Murphy [38] proposed as well a family of 8 parsimonious Gaussian models by extending the mixture of factor analyzers (MFA) model [37, 42] by constraining the loading and error variance matrices across groups. Let us remark that the two families of parsimonious Gaussian models share some models: for instance, the model UUC of [38] corresponds to the model  $[a_{kj}b_kQ_kd]$  of [9]. Among the 8 parsimonious models presented in [38], 4 models have the loading matrices constrained across the groups. More recently, Beak *et al.* [5] proposed as well a MFA model with a common loading matrix. All these parsimonious Gaussian models share the assumption that the group subspaces have a common orientation and are therefore close to the DLM models presented in this work. However, these models with common loadings choose the orientation such as the variance of the projected data is maximum whereas the DLM models choose the latent subspace orientation such as it best discriminates the groups. This specific feature of the DLM models should therefore improve in most cases both the clustering and the visualization of the results.

## 4 Parameter estimation: the Fisher-EM algorithm

This section introduces an estimation procedure which adapts the traditional EM algorithm for estimating the parameters of DLM models presented in the previous section. Due to the nature of the models described above, the Fisher-EM algorithm alternates between three steps:

- an E step in which posterior probabilities that observations belong to the  $K$  groups are computed,
- a F step which estimates the orientation matrix  $U$  of the discriminative latent space conditionally to the posterior probabilities,
- a M step in which parameters of the mixture model are estimated in the latent subspace by maximizing the conditional expectation of the complete likelihood.

This estimation procedure relative to the DLM models is called hereafter the Fisher-EM algorithm. We chose to name this estimation procedure after Sir R. A. Fisher since the key-idea of the F step comes from his famous work on discrimination. The remainder of this section details the simple form of this procedure. Let us however notice that the Fisher-EM algorithm can be also used in combination with the stochastic [10] and classification versions [11] of the EM algorithm.

### 4.1 The E step

This step aims to compute, at iteration  $(q)$ , the expectation of the complete log-likelihood conditionally to the current value of the parameter  $\theta^{(q-1)}$ , which, in practice, resumes to the computation of  $t_{ik}^{(q)} = E[z_{ik}|y_i, \theta^{(q-1)}]$ . Let us also recall that  $t_{ik}^{(q)}$  is as well the posterior probability that the observation  $y_i$  belongs to the  $k^{th}$  component of the mixture. The following proposition provides the explicit form of  $t_{ik}^{(q)}$ , for  $i = 1, \dots, n$ ,  $k = 1, \dots, K$ . Demonstration of this result is detailed in Appendix A.1.

**Proposition 1.** *With the assumptions of the DLM models, the posterior probabilities  $t_{ik}^{(q)}$ ,  $i = 1, \dots, n$ ,  $k = 1, \dots, K$ , can be expressed as :*

$$t_{ik}^{(q)} = \frac{1}{\sum_{l=1}^K \exp\left(\frac{1}{2}(\Gamma_k^{(q-1)}(y) - \Gamma_l^{(q-1)}(y))\right)},$$

with:

$$\begin{aligned} \Gamma_k^{(q-1)}(y_i) = & \|P(y_i - m_k^{(q-1)})\|_{\mathcal{D}_k}^2 + \frac{1}{\beta_k^{(q-1)}} \|(y_i - m_k^{(q-1)}) - P(y_i - m_k^{(q-1)})\|^2 \\ & + \sum_{j=1}^d \log(\alpha_{kj}^{(q-1)}) + (p-d) \log(\beta_k^{(q-1)}) - 2 \log(\pi_k^{(q-1)}) + \gamma, \end{aligned} \quad (4.1)$$




Figure 2: Two groups and their 1-dimensional discriminative subspace.

where  $\|\cdot\|_{\mathcal{D}_k}^2$  is a norm on the latent space  $\mathbb{E}$  defined by  $\|y\|_{\mathcal{D}_k}^2 = y^T \mathcal{D}_k y$ ,  $\mathcal{D}_k = \tilde{W} \Delta_k^{-1} \tilde{W}^T$ ,  $\tilde{W}$  is a  $p \times p$  matrix containing the  $d$  vectors of  $U^{(q-1)}$  completed by zeros such as  $\tilde{W} = [U^{(q-1)}, 0_{p-d}]$ ,  $P$  is the projection operator on the latent space  $\mathbb{E}$ , i.e.  $P(y) = U^{(q-1)} U^{(q-1)T} y$ , and  $\gamma = p \log(2\pi)$  is a constant term.

Besides its computational interest, Proposition 1 provides as well a comprehensive interpretation of the cost function  $\Gamma_k$  which mainly governs the computation of  $t_{ik}$ . Indeed, it appears that  $\Gamma_k$  mainly depends on two distances: the distance between the projections on the discriminant subspace  $\mathbb{E}$  of the observation  $y_i$  and the mean  $m_k$  on the one hand, and, the distance between the projections on the complementary subspace  $\mathbb{E}^\perp$  of  $y_i$  and  $m_k$  on the other hand. Remark that the latter distance can be reformulated in order to avoid the use of the projection on  $\mathbb{E}^\perp$ . Indeed, as Figure 2 illustrates, this distance can be re-expressed according projections on  $\mathbb{E}$ . Therefore, the posterior probability  $t_{ik} = P(z_{ik} = 1 | y_i)$  will be close to 1 if both the distances are small which seems quite natural. Obviously, these distances are also balanced by the variances in  $\mathbb{E}$  and  $\mathbb{E}^\perp$  and by the mixture proportions. Furthermore, the fact that the E step does not require the use of the projection on the complementary subspace  $\mathbb{E}^\perp$  is, from a computational point of view, very important because it will provide the stability of the algorithm and will allow its use when  $n \ll p$  (cf. paragraph 4.5).

## 4.2 The F step

This step aims to determinate, at iteration  $(q)$ , the discriminative latent subspace of dimension  $d = K - 1$  in which the  $K$  groups are best separated. Naturally, the estimation of this latent subspace has to be done conditionally to the current values of posterior probabilities  $t_{ik}^{(q)}$  which indicates the current fuzzy partition of the data. Estimating the discriminative latent subspace  $\mathbb{E}^{(q)}$  resumes to the computation of  $d$  discriminative axes. Following the original idea of Fisher [15], the  $d$  axes which best discriminate the  $K$  groups are those which maximize the traditional criterion  $J(U) = \text{tr}((U^T S_B U)^{-1} U^T S_W U)$ . However, the traditional criterion  $J(U)$  assume that the data are complete (supervised classification framework). Unfortunately, the situation of interest here is the one of the unsupervised classification and the matrices  $S_B$  and  $S_W$  have therefore to be defined conditionally to the current fuzzy partition. Furthermore, the DLM models assume that the discriminative latent subspace must have an orthonormal basis and, sadly, the traditional Fisher's approach provides correlated discriminative axes.

To overcome both problems, this paragraph proposes a procedure which keeps the key-idea of Fisher while providing orthonormal discriminative axes conditionally to the current fuzzy partition of the data. The procedure follows the concept of the orthonormal discriminant vector (ODV) method introduced by [16] in the supervised case and then extended by [21, 22, 33, 50], which sequentially selects the most discriminative features in maximizing the Fisher criterion subject to the orthogonality of features. First, it is necessary to introduce the fuzzy between-covariance matrix  $S_B^{(q)}$  and the fuzzy within-covariance matrix  $S_W^{(q)}$ . The fuzzy between-covariance matrix  $S_B^{(q)}$  is defined conditionally to the posterior probabilities  $t_{ik}^{(q)}$ , obtained in the E step, as follows:

$$S_B^{(q)} = \frac{1}{n} \sum_{k=1}^K n_k^{(q)} (\hat{m}_k^{(q)} - \bar{y})^T (\hat{m}_k^{(q)} - \bar{y}),$$

where  $n_k^{(q)} = \sum_{i=1}^n t_{ik}^{(q)}$ ,  $\hat{m}_k^{(q)} = \frac{1}{n} \sum_{i=1}^n t_{ik}^{(q)} y_i$  is the fuzzy mean of the  $k$ th group at iteration  $q$  and  $\bar{y} = \frac{1}{n} \sum_{i=1}^n y_i$  is the empirical mean of the whole dataset. Since the relation  $S = S_W^{(q)} + S_B^{(q)}$  holds in this context as well, it is preferable from a computational point of view to use the covariance matrix  $S = \frac{1}{n} \sum_{i=1}^n (y_i - \bar{y})(y_i - \bar{y})^T$  of the whole dataset in the maximization problem instead  $S_W^{(q)}$  since  $S$  remains fixed over the iteration. The F step of the Fisher-EM therefore aims to solve the following optimization problem:

$$\begin{cases} \max_U & \text{tr} \left( (U^T S U)^{-1} U^T S_B^{(q)} U \right), \\ \text{wrt} & u_j^T u_l = 0, \quad \forall j \neq l \in \{1, \dots, d\}, \end{cases}$$

where  $u_j$  is the  $j$ th column vector of  $U$ . Following the ODV procedure, the  $d$  axes solution of this optimization problem are iteratively constructed by, first, computing an orthogonal complementary subspace to the current set of discriminative axes and, then, maximizing the

Fisher criterion in this orthogonal subspace. To initialize this iterative procedure, the first vector of  $U$  is obtained using the traditional maximization of the Fisher criterion and  $u_1$  is therefore the eigenvector associated with the largest eigenvalue of the matrix  $S^{-1}S_B^{(q)}$ . Then, assuming that the  $r - 1$  first orthonormal discriminative axes  $\{u_1, \dots, u_{r-1}\}$ , which span the space  $\mathcal{B}^{r-1}$ , have been computed, the  $r^{th}$  discriminative axis has to lie in the subspace  $\mathcal{B}^r$  orthogonal to the space  $\mathcal{B}^{r-1}$ . The Gram-Schmidt orthonormalization procedure allows to find a basis  $V^r = \{v_r, v_{r+1}, \dots, v_d\}$  for the orthogonal subspace  $\mathcal{B}^r$  such that:

$$v_\ell = \alpha_\ell (I_{\ell-1} - \sum_{j=1}^{\ell-1} v_j v_j^T) \psi_\ell, \quad \ell = r, \dots, p$$

where  $v_j = u_j$  for  $j = 1, \dots, r - 1$ ,  $\alpha_\ell$  is normalization constant such that  $\|u_\ell\| = 1$  and  $\psi_\ell$  is a vector linearly independent of  $u_j \forall j \in \{1, \dots, \ell - 1\}$ . Then, the  $r$ th discriminative axis is given by:

$$u_r = \frac{P_{r-1} u_r^{max}}{\|u_r^{max}\|},$$

where  $P_{r-1}$  is the projector on  $\mathcal{B}^{r-1}$ ,  $u_r^{max}$  is the eigenvector associated with the largest eigenvalue of the matrix  $S_r^{-1}S_{B^r}^{(q)}$  with:

$$\begin{aligned} S_r &= V^{rT} S V^r, \\ S_{B^r}^{(q)} &= V^{rT} S_B^{(q)} V^r, \end{aligned}$$

i.e.  $S_r$  and  $S_{B^r}^{(q)}$  are respectively the covariance and fuzzy between-covariance matrices of the data projected into the orthogonal subspace  $\mathcal{B}^r$ . This iterative procedure stops when the  $d = K - 1$  orthonormal discriminative axes  $u_j$  are computed.

### 4.3 The M step

This third step estimates the model parameters by maximizing the conditional expectation of the complete likelihood. The following proposition provides the expression of the conditional expectation of the complete log-likelihood in the case of the  $\text{DLM}_{[\alpha_{kj}\beta_k]}$  model. Demonstration of this result is provided in Appendix A.2

**Proposition 2.** *In the case of the  $\text{DLM}_{[\alpha_{kj}\beta_k]}$  model, the conditional expectation of complete log-likelihood  $Q(y_1, \dots, y_n, \theta)$  has the following expression:*

$$\begin{aligned} Q(y_1, \dots, y_n, \theta) = & -\frac{1}{2} \sum_{k=1}^K n_k \left[ -2 \log(\pi_k) + \sum_{j=1}^d \left( \log(\alpha_{kj}) + \frac{u_j^T C_k u_j}{\alpha_{jk}} \right) \right. \\ & \left. + (p - d) \log(\beta_k) + \frac{1}{\beta_k} \sum_{j=d+1}^p v_j^T C_k v_j + \gamma \right]. \end{aligned} \quad (4.2)$$



where  $C_k$  is the empirical covariance matrix of the  $k^{th}$  group,  $u_j$  is the  $j$ th column vector of  $U$ ,  $v_j$  is the  $j$ th column vector of  $V$ ,  $n_k = \sum_{i=1}^n t_{ik}$  and  $\gamma = p \log(2\pi)$  is a constant term.

At iteration  $q$ , the maximization of  $Q$  conduces to an estimation of the mixture proportions  $\pi_k$  and the means  $\mu_k$  for the  $K$  components by their empirical counterparts:

$$\begin{aligned}\hat{\pi}_k^{(q)} &= \frac{n_k}{n}, \\ \hat{\mu}_k^{(q)} &= \frac{1}{n_k} \sum_{i=1}^n t_{ik}^{(q)} U^{(q)T} y_i,\end{aligned}$$

where  $n_k = \sum_{i=1}^n t_{ik}^{(q)}$  and  $U^{(q)}$  contains, as columns vectors, the  $d$  discriminative axes  $u_j^{(q)}$ ,  $j = 1, \dots, d$ , provided by the F step at iteration  $q$ . The following proposition provides estimates for the remaining parameters for the 8 DLM models which have to be updated at each iteration of the FEM procedure. Demonstrations of the following results are given in Appendix A.2.

**Proposition 3.** *At iteration  $q$ , the estimates for variance parameters of the 8 DLM models are:*

- *Model DLM $_{[\alpha_{kj}\beta_k]}$ :*

$$\hat{\alpha}_{kj}^{(q)} = u_j^{(q)T} C_k^{(q)} u_j^{(q)}, \quad (4.3)$$

$$\hat{\beta}_k^{(q)} = \frac{\text{tr}(C_k^{(q)}) - \sum_{j=1}^d u_j^{(q)T} C_k^{(q)} u_j^{(q)}}{p - d}, \quad (4.4)$$

- *Model DLM $_{[\alpha_{kj}\beta]}$ :*

$$\hat{\alpha}_{kj}^{(q)} = u_j^{(q)T} C_k^{(q)} u_j^{(q)}, \quad (4.5)$$

$$\hat{\beta}^{(q)} = \frac{\text{tr}(C^{(q)}) - \sum_{j=1}^d u_j^{(q)T} C^{(q)} u_j^{(q)}}{p - d}, \quad (4.6)$$

- *Model DLM $_{[\alpha_k\beta_k]}$ :*

$$\hat{\alpha}_k^{(q)} = \frac{1}{d} \sum_{j=1}^d u_j^{(q)T} C_k^{(q)} u_j^{(q)}, \quad (4.7)$$

$$\hat{\beta}_k^{(q)} = \frac{\text{tr}(C_k^{(q)}) - \sum_{j=1}^d u_j^{(q)T} C_k^{(q)} u_j^{(q)}}{p - d}, \quad (4.8)$$

- *Model DLM*<sub>[ $\alpha_k \beta$ ]</sub>:

$$\hat{\alpha}_k^{(q)} = \frac{1}{d} \sum_{j=1}^d u_j^{(q)T} C_k^{(q)} u_j^{(q)}, \quad (4.9)$$

$$\hat{\beta}^{(q)} = \frac{\text{tr}(C^{(q)}) - \sum_{j=1}^d u_j^{(q)T} C^{(q)} u_j^{(q)}}{p - d}, \quad (4.10)$$

- *Model DLM*<sub>[ $\alpha_j \beta_k$ ]</sub>:

$$\hat{\alpha}_j^{(q)} = u_j^{(q)T} C^{(q)} u_j^{(q)}, \quad (4.11)$$

$$\hat{\beta}_k^{(q)} = \frac{\text{tr}(C_k^{(q)}) - \sum_{j=1}^d u_j^{(q)T} C_k^{(q)} u_j^{(q)}}{p - d}, \quad (4.12)$$

- *Model DLM*<sub>[ $\alpha_j \beta$ ]</sub>:

$$\hat{\alpha}_j^{(q)} = u_j^{(q)T} C^{(q)} u_j^{(q)}, \quad (4.13)$$

$$\hat{\beta}^{(q)} = \frac{\text{tr}(C^{(q)}) - \sum_{j=1}^d u_j^{(q)T} C^{(q)} u_j^{(q)}}{p - d}, \quad (4.14)$$

- *Model DLM*<sub>[ $\alpha \beta_k$ ]</sub>:

$$\hat{\alpha}^{(q)} = \frac{1}{d} \sum_{j=1}^d u_j^{(q)T} C^{(q)} u_j^{(q)}, \quad (4.15)$$

$$\hat{\beta}_k^{(q)} = \frac{\text{tr}(C_k^{(q)}) - \sum_{j=1}^d u_j^{(q)T} C_k^{(q)} u_j^{(q)}}{p - d}, \quad (4.16)$$

- *Model DLM*<sub>[ $\alpha \beta$ ]</sub>:

$$\hat{\alpha}^{(q)} = \frac{1}{d} \sum_{j=1}^d u_j^{(q)T} C^{(q)} u_j^{(q)}, \quad (4.17)$$

$$\hat{\beta}^{(q)} = \frac{\text{tr}(C^{(q)}) - \sum_{j=1}^d u_j^{(q)T} C^{(q)} u_j^{(q)}}{p - d}, \quad (4.18)$$

where the vectors  $u_j^{(q)}$  are the discriminative axes provided by the  $F$  step at iteration  $q$ ,  $C_k^{(q)} = \frac{1}{n_k^{(q)}} \sum_{i=1}^n t_{ik}^{(q)} (y_i - \hat{m}_k^{(q)})^T (y_i - \hat{m}_k^{(q)})$  is the fuzzy covariance matrix of the  $k$ th group,  $\hat{m}_k^{(q)} = \frac{1}{n} \sum_{i=1}^n t_{ik}^{(q)} y_i$  and finally  $C = \frac{1}{n} \sum_{k=1}^K n_k C_k$  is the fuzzy within-covariance matrix of the  $K$  groups.

#### 4.4 Initialization and model selection

Since the Fisher-EM procedure presented in this work belongs to the family of EM-based algorithms, the Fisher-EM algorithm can inherit the most efficient strategies for initialization and model selection from previous works on the EM algorithm.

**Initialization** Although the EM algorithm is widely used, it is also well-known that the performance of the algorithm is linked to its initial conditions. Several strategies have been proposed in the literature for initializing the EM algorithm. A popular practice [8] executes the EM algorithm several times from a random initialization and keep only the set of parameters associated with the highest likelihood. The use of kmeans or of a random partition are also standard approaches for initializing the algorithm. McLachlan and Peel [36] have also proposed an initialization through the parameters by generating the mean and the covariance matrix of each mixture component from a multivariate normal distribution parametrized by the empirical mean and empirical covariance matrix of the data. In practice, this latter initialization procedure works well but, unfortunately, it can not be applied directly to the Fisher-EM algorithm since model parameters live in a space different from the observation space. A simple way to adapt this strategy could be to first determine a latent space using PCA and then simulate mixture parameters in this initialization latent space.

**Model selection** In model-based clustering, it is frequent to consider several models in order to find the most appropriate model for the considered data. Since a model is defined by its number of component  $K$  and its parametrization, model selection allows to both select a parametrization and a number of components. Several criteria for model selection have been proposed in the literature and the famous ones are penalized likelihood criteria. Classical tools for model selection include the AIC [2], BIC [44] and ICL [7] criteria. The Bayesian Information Criterion (BIC) is certainly the most popular and consists in selecting the model which penalizes the likelihood by  $\frac{\gamma(\mathcal{M})}{2} \log(n)$  where  $\gamma(\mathcal{M})$  is the the number of parameters in model  $\mathcal{M}$  and  $n$  is the number of observations. On the other hand, the AIC criterion penalizes the log-likelihood by  $\gamma(\mathcal{M})$  whereas the ICL criterion add the penalty  $\sum_{i=1}^n \sum_{k=1}^K t_{ik} \log(t_{ik})$  to the one of the BIC criterion in order to favor well separated models. The value of  $\gamma(\mathcal{M})$  is of course specific to the model selected by the practician (*cf.* Table 1). In the experiments of the following sections, the BIC criterion is used because of its popularity but the ICL criterion should also be well adapted in our context.

#### 4.5 Practical and numerical considerations

The DLM models, for which the Fisher-EM algorithm has been proposed as an estimation procedure, presents several practical and numerical interests among which the ability to deal with the so-called  $n \ll p$  problem and to interpret the discriminative axes.

**Use and interpretation of the discriminative axes** The first and natural use of the discriminative axes may certainly be the visualization of the clustered data. Indeed, it is nowadays clear that the visualization help human operators to understand the results of an analysis. With the Fisher-EM algorithm, it is easy to project and visualize the cluster data into the estimated discriminative latent subspace. Beyond this natural interest, it may be useful from a practical point of view to interpret the  $d = K - 1$  estimated discriminative axes, *i.e.*  $u_1, \dots, u_d$  with the notations of the previous sections. The main interest for the practitioner would be to figure out which original dimensions are the most discriminative. This can be done by looking at the matrix  $U$  which contains  $u_1, \dots, u_d$  as column vectors. In the classical framework of factor analysis, this matrix is known as the loading matrix (the discriminative axes  $u_1, \dots, u_d$  are the loadings). The loadings link the discriminative latent space with the original observation space and allow to calculate the correlation between the discriminative axes and the original variables. Indeed, the correlation between the  $j$ th discriminative axes  $u_j$ ,  $j = 1, \dots, d$ , and the  $\ell$ th original variable  $v_\ell$  is given by  $c_{j\ell} = \sqrt{\lambda_j} u_{j\ell}$  where  $\lambda_j$  is the eigenvalue associated with  $u_j$ . Thus, it is possible to find the most discriminative original variables by selecting the the highest correlated original variables with the discriminative axes. Let us finally remark that finding the most discriminative original variables is of particular interest in application fields, such as biology or economics, where the observed variables have an actual meaning.

**Dealing with the  $n \ll p$  problem** Another important and frequent problem when clustering high-dimensional data is known as high dimension and low sample size (HDSS) problem or the  $n \ll p$  problem (we refer to [25, Chap. 18] for an overview). The  $n \ll p$  problem refers to situations where the number of features  $p$  is larger than the number of available observations  $n$ . This problem occurs frequently in modern scientific applications such as genomics or mass spectrometry. In such cases, the estimation of model parameters for generative clustering methods is either difficult or impossible. This task is indeed very difficult when  $n \ll p$  since generative methods require, in particular, to invert covariance matrices which are ill-conditioned in the best case or singular in the worst one. In contrast with other generative methods, the Fisher-EM procedure can overcome the  $n \ll p$  problem. Indeed, the E and M steps of Fisher-EM do not require the determination of the last  $p - d$  columns of  $W$  (see equations (4.1) and (4.3)–(4.18)) and, consequently, it is possible to modify the F step to deal with situations where  $n \ll p$ . To do so, let  $\bar{Y}$  denote the centered data matrix and  $T$  denote, as before, the fuzzy partition matrix. We define in addition the weighted fuzzy partition matrix  $\tilde{T}$  where the  $j$ th column  $\tilde{T}_j$  of  $\tilde{T}$  is the  $j$ th column  $T_j$  of  $T$  divided by  $n_j = \sum_{i=1}^n t_{ij}$ . With these notations, the between covariance matrix  $B$  can be written in its matrix form  $B = \bar{Y}^t \tilde{T}^t \tilde{T} \bar{Y}$  and the F step aims to maximize, under orthogonality constraints, the following function:

$$f(U) = \text{tr} \left( (U^t \bar{Y}^t \bar{Y} U)^{-1} U^t \bar{Y}^t \tilde{T}^t \tilde{T} \bar{Y} U \right). \quad (4.19)$$

It follows from the classical result of kernel theory, the Representer theorem, that this maximization can be done in a different space and that  $U$  can be expressed as  $U = \bar{Y}H$  where  $H \in \mathbb{R}^{n \times p}$ . Therefore, the F step reduces to maximize, under orthogonality constraints, the following function:

$$f(H) = \text{tr} \left( (H^t G G H)^{-1} H^t G \tilde{T}^t \tilde{T} G H \right), \quad (4.20)$$

where  $G = \bar{Y} \bar{Y}^t$  is the  $n \times n$  Gram matrix. The solution  $U^*$  of (4.19) can be obtained afterward from the solution  $H^*$  of (4.20) by multiplying it by  $\bar{Y}$ . Thus, the F step reduces to the eigendecomposition under orthogonality constraints of a  $n \times n$  matrix instead of a  $p \times p$  matrix. This procedure is useful for the Fisher-EM procedure only because it allows to determine  $d \leq n$  axes which are enough for Fisher-EM but not for other generative methods which require the computation of the  $p$  axes.

## 5 Experimental results

This section presents experiments on simulated and real datasets in order to highlight the main features of the clustering method introduced in the previous sections.

### 5.1 An introductory example: the Fisher’s irises

Since we chose to name the clustering algorithm proposed in this work after Sir R. A. Fisher, the least we can do is to first apply the Fisher-EM algorithm to the iris dataset that Fisher used in [15] as an illustration for his discriminant analysis. This dataset, in fact collected by E. Anderson [4] in the Gaspé peninsula (Canada), is made of three groups corresponding to different species of iris (*setosa*, *versicolor* and *virginica*) among which the groups *versicolor* and *virginica* are difficult to discriminate (they are at least not linearly separable). The dataset consists of 50 samples from each of three species and four features were measured from each sample. The four measurements are the length and the width of the sepal and the petal. This dataset is used here as an introductory example because of the link with Fisher’s work but also of its popularity in the clustering community.

In this first experiment, Fisher-EM has been applied to the iris data (of course, the labels have been used only for performance evaluation) and the Fisher-EM results will be compared to the ones obtained in the supervised case with the orthogonal linear analysis method (OLDA) [50]. The left panel of Figure 3 stands for the projection of the irises in the estimated discriminative space with Fisher-EM and the right panel shows the evolution of the log-likelihood on 25 iterations until convergence. First of all, it can be observed that the estimated latent space discriminates almost perfectly the three different groups. For this experiment, the clustering accuracy has reached 98% with the  $\text{DLM}_{[\alpha_k, \beta]}$  model of Fisher-EM. Secondly, the right panel shows the monotony of the evolution of the log-likelihood and the convergence of the algorithm to a stationary state. Table 2 presents the confusion matrices for




Figure 3: Projection of clustered Iris data into the latent discriminative subspace with Fisher-EM (left) and evolution of the associated log-likelihood (right).

OLDA				Fisher-EM			
<i>cluster</i>				<i>cluster</i>			
<i>class</i>	1	2	3	<i>class</i>	1	2	3
Setosa	50	0	0	Setosa	50	0	0
Versicolor	0	48	2	Versicolor	0	47	3
Virginica	0	1	49	Virginica	0	0	50
<i>Misclassification rate = 0.02</i>				<i>Misclassification rate = 0.02</i>			

Table 2: Confusion tables for the iris data with OLDA method (supervised) and Fisher-EM (unsupervised).

<i>variable</i>	OLDA		Fisher-EM	
	<i>axis</i>		<i>axis</i>	
	1	2	1	2
sepal length	0.209	0.044	-0.203	-0.108
sepal width	0.386	0.665	-0.422	0.088
petal length	-0.554	-0.356	0.602	0.736
petal width	-0.707	0.655	0.646	-0.662

Table 3: Fisher axes estimated in the supervised case (OLDA) and in the unsupervised case (Fisher-EM).




Figure 4: Visualization of the simulated data: data in their latent space (left) and data projected on the first principal components (right).

the partitions obtained with supervised and unsupervised classification methods. OLDA has been used for the supervised case (reclassification of the learning data) whereas Fisher-EM has provided the clustering results. One can observe that the obtained partitions induced by both methods is almost the same. This confirms that Fisher-EM has correctly modeled both the discriminative subspace and the groups within the subspace. It is also interesting to look at the loadings provided by both methods. Table 3 stands for the linear coefficients of the discriminative axes estimated, on the one hand, in the supervised case (OLDA) and, on the other hand, in the unsupervised case (Fisher-EM). The first axes of each approach appear to be very similar and the scalar product of these axes is  $-0.996$ . This highlights the performance of the Fisher-EM algorithm in estimating the discriminative subspace of the data. Furthermore, according to these results, the 3 groups of irises can be mainly discriminated by the petal size meaning that only one axis would be sufficient to discriminate the 3 iris species. Besides, this interpretation turns out to be in accordance with the recent work of Trendafilov and Joliffe [47] on variable selection in discriminant analysis via the LASSO.

## 5.2 Simulation study: influence of the dimension

This second experiment aims to compare with traditional methods the stability and the efficiency of the Fisher-EM algorithm in partitioning high dimensional data. Fisher-EM is compared here with the standard EM algorithm (Full-GMM) and its parsimonious models (Diag-GMM, Sphe-GMM and Com-GMM), the EM algorithm applied in the first components of PCA explaining 90% of the total variance (PCA-EM), the kmeans algorithm and the mixture of probabilistic principal component analyzers (Mixt-PPCA). For this simulation, 600




Figure 5: Influence of the dimension of the observed space on the correct classification rate for Full-GMM, PCA-EM, Com-GMM, Mixt-PPCA, kmeans, Diag-GMM, Sphe-GMM and Fisher-EM algorithms.

observations have been simulated following the  $DLM_{[\alpha_{kj}\beta_k]}$  model proposed in Section 3. The simulated dataset is made of 3 unbalanced groups and each group is modeled by a Gaussian density in a 2-dimensional space completed by orthogonal dimensions of Gaussian noise. The transformation matrix  $W$  has been randomly simulated such as  $W^T W = W W^T = I_p$  and, for this experience, the dimension of the observed space varies from 5 to 100. The left panel of Figure 4 shows the simulated data in their 2-dimensional latent space whereas the right panel presents the projection of 50-dimensional observed data on the two first axes of PCA in the observed space. As one can observe, the representation of the data on the two first principal components is actually not well suited for clustering these data while it exists a representation which discriminates perfectly the three groups. Moreover, to make the results of each method comparable, the same randomized initialization has been used for the 8 algorithms. The experimental process has been repeated 20 times for each dimension of the observed space in order to see both the average performances and their variances. Figure 5 presents the evolution of the clustering accuracy of each method (EM, PCA-EM, kmeans, Mixt-PPCA, Fisher-EM, Diag-GMM, Sphe-GMM and Com-GMM) according to the data dimensionality and Figure 6 presents their respective boxplots. First of all, it can be observed that the Full-GMM, PCA-EM and Com-GMM have their performances which decrease quickly when the dimension increases. In fact, the Full-GMM model does not work upon the 15th dimension and still remains unstable in a low dimensional space as well as the Com-GMM model. Sim-






Figure 6: Boxplots of Full-GMM, PCA-EM, Com-GMM, Mixt-PPCA, kmeans, Diag-GMM, Sphe-GMM and Fisher-EM algorithms.

ilarly, the performances of PCA-EM fall down as the 10th dimension. This can be explained by the fact that the latent subspace provided by PCA does not allow to well discriminate the groups, as already suggested by Figure 4. However, the PCA-EM approach can be used whatever the dimension is whereas Full-GMM can not be used as the 20th dimension because of numerical problems linked to singularity of the covariance matrices. Moreover, their box-plots show a large variation on the clustering accuracy. Secondly, Sphe-GMM, Diag-GMM and kmeans present the same trend with high performances in low-dimensional spaces which decrease until they reach a clustering accuracy of 0.75. Diag-GMM seems however to resist a little bit more than kmeans to the dimension increasing. Mixt-PPCA and Mclust both follow the same tendency as the previous methods but from the 30th dimension their performances fall down until the clustering accuracy reaches 0.50. The poor performances of Mixt-PPCA can be explained by the fact that Mixt-PPCA models each group in a different subspace whereas the model used for simulating the observations assumes a common discriminative subspace. Finally, Fisher-EM appears to be more efficient than the other methods and, more importantly, it remains very stable while the data dimensionality increases. Furthermore, the boxplot associated with the Fisher-EM results suggests that it is a steady algorithm which succeeds in finding out the discriminative latent subspace of the data even with random initializations.

### 5.3 Simulation study: BIC criterion

This last experiment on simulations aims to study the performance of BIC for both model and component number selection. For this experiment, 4 Gaussian components have been simulated according to the  $DLM_{[\alpha_k, \beta]}$  model in a 3-dimensional space completed by 17 orthogonal dimensions of Gaussian noise. The transformation matrix  $W$  has been again randomly simulated such as  $W^T W = W W^T = I_p$ . Table 4 presents the BIC values for the family of DLM models and, in a comparative purpose, the BIC values for 7 other methods already used in the last experiments: EM with the Full-GMM, Diag-GMM, Sphe-GMM and Com-GMM models, Mixt-PPCA, Mclust (with its most adapted model for these data) and PCA-EM. Moreover, BIC is computed for different partition numbers varying between 2 and 6 clusters. First of all, one can observe that the BIC values linked to the models which are different from the DLM model are very low compared to the DLM models. This suggests that the models which best fit the data are the DLM models. Secondly, only some DLM models, those which assume a different variance between each cluster such as  $DLM_{\alpha_{kj}\beta_k}$ ,  $DLM_{\alpha_{kj}\beta}$ ,  $DLM_{\alpha_k\beta_k}$  and  $DLM_{\alpha_k\beta}$ , select the right number of components ( $K = 4$ ). The other methods either over-estimate the number of clusters, such as Mclust, or under-estimate it. BIC has the smaller value for the  $DLM_{\alpha_k\beta}$  model with 4 components which is actually the model used for simulating the data. Finally, the right-hand side of Figure 7 presents the projection of the data on the discriminant subspace of 3 dimensions estimated by Fisher-EM with the  $DLM_{\alpha_k\beta}$  model whereas the left-hand side figure represents the projection of the data on the 3 first principal components

of PCA. As one can observe, in the PCA case, the axes separate only 2 groups, which is in accordance with the model selection pointed out by BIC for this method. Conversely, in the Fisher-EM case, the 3 discriminative axes separate well the 4 groups and such a representation could clearly help the practitioner in understanding the clustering results.

#### 5.4 Real data set benchmark

This last experimental paragraph will focus on comparing on real-world datasets the efficiency of Fisher-EM with several linear and nonlinear existing methods, including the most recent ones. On the one hand, Fisher-EM will be compared to the 8 already used clustering methods: EM with the Full-GMM, Diag-GMM, Sphe-GMM and Com-GMM models, Mixt-PPCA, Mclust (with its most adapted model for these data), PCA-EM and kmeans. On the other hand, the new Fisher-EM challengers will be kmeans computed on the two first components of PCA (PCA-kmeans), an heteroscedastic factor mixture analyzer (HMFA) method [39] and three discriminative versions of kmeans: LDA-kmeans [13], Diskmeans and DisCluster (see [51] for more details). The comparison has been made on 6 different benchmark datasets coming mostly from the UCI machine learning repository:


- The **chironomus** data contain 148 larvae which are split up into 3 species and described by 17 morphometric attributes. This dataset is described in detailed in [39].
- The **wine** dataset is composed by 178 observations which are split up into 3 classes and characterized by 13 variables.
- The **iris** dataset which is made of 3 different groups and described by 4 variables. This dataset has been described in detail in Section 5.1.
- The **zoo** dataset includes 7 families of 101 animals characterized by 16 variables.
- The **glass** data are composed by 214 observations belonging to 6 different groups and described by 7 variables.
- Finally, the 4435 **satellite images** are split up into 6 classes and are described by 36 variables.

Table 5 presents the average clustering accuracies obtained for the 8 DLM models and for the methods already used in the previous experiments. The results for the 14 first methods of the table have been obtained by averaging 20 trials with random initializations. Similarly, Table 6 provides the clustering accuracies found in the literature for the recent methods on the same datasets. It is important to notice that the results of Table 6 have been obtained in slightly different benchmarking situations. Missing values in Table 5 are due to non convergence of the algorithms whereas missing values in Table 6 are due to the unavailability of the information for the concerned method. First of all, one can remark that Fisher-EM outperforms the

methods	number of components				
	2	3	4	5	6
$\text{DLM}_{\alpha_k j \beta_k}$	-83.419	-83.231	-82.967	-83.404	-83.352
$\text{DLM}_{\alpha_k j \beta}$	-83.750	-83.275	-82.971	-83.398	-83.885
$\text{DLM}_{\alpha_k \beta_k}$	-83.419	-83.239	-82.970	-83.596	-84.136
$\text{DLM}_{\alpha_k \beta}$	-83.750	-83.536	<b>-82.956</b>	-83.616	-84.082
$\text{DLM}_{\alpha_j \beta_k}$	-84.782	-85.219	-85.557	-86.113	-86.416
$\text{DLM}_{\alpha_j \beta}$	-84.764	-85.189	-85.668	-86.208	-86.230
$\text{DLM}_{\alpha \beta_k}$	-84.782	-85.313	-85.914	-86.309	-86.607
$\text{DLM}_{\alpha \beta}$	-84.764	-85.467	-85.910	-86.516	-86.793
Full-GMM	-5012.136	-7532.103	-10063.559	-12628.357	-15143.680
Com-GMM	-1866.557	-2091.528	-2317.083	-2543.283	-2770.033
Mixt-PPCA	-4932.571	-7319.903	-9707.076	-12093.666	-14481.251
Diag-GMM	-940.067	-1418.939	-1902.644	-2384.809	-2873.094
Sphe-GMM	-613.337	-931.568	-1244.007	-1564.471	-1885.092
PCA-EM	-3341.966	-5024.925	-6715.415	-8416.745	-10115.715
Mclust <sub>[EEE]</sub>	-166.956	-167.083	-167.180	-167.366	-165.962

Table 4: BIC values for model selection.

Projection on the 3 first principal components



Projection on the discriminative axes estimated by Fisher-EM




Figure 7: Projection of the data in the 3 first principal components of PCA (left) and in the discriminant subspace estimated by Fisher-EM with the  $\text{DLM}_{[\alpha_k \beta]}$ .

Method	chironomus	wine	iris	zoo	glass	satimage
DLM $_{\alpha_k j \beta_k}$	0.860	0.955	0.893	0.682	<b>0.439</b>	<b>0.665</b>
DLM $_{\alpha_k j \beta}$	<b>0.987</b>	0.942	0.906	0.782	0.387	0.657
DLM $_{\alpha_k \beta_k}$	0.887	0.955	0.960	0.752	-	0.628
DLM $_{\alpha_k \beta}$	<b>0.987</b>	0.943	<b>0.980</b>	0.697	0.372	0.649
DLM $_{\alpha_j \beta_k}$	0.836	0.942	0.827	0.717	0.320	0.621
DLM $_{\alpha_j \beta}$	0.897	0.932	0.667	<b>0.802</b>	0.385	0.662
DLM $_{\alpha \beta_k}$	0.836	0.954	0.793	0.718	0.320	0.552
DLM $_{\alpha \beta}$	0.897	0.966	0.800	0.776	0.385	0.586
Full-GMM	0.445	0.911	0.766	-	0.377	0.486
Sphe-GMM	0.760	0.721	0.894	0.732	0.373	0.629
Diag-GMM	0.899	0.943	0.854	-	-	0.647
Comm-GMM	0.548	0.725	0.530	-	0.403	0.266
Mixt-PPCA	0.604	0.494	0.688	0.512	0.385	0.438
Mclust ( <i>Model name</i> )	0.979 ( <i>EEE</i> )	<b>0.971</b> ( <i>VVI</i> )	0.966 ( <i>VEV</i> )	0.584 ( <i>EII</i> )	0.415 ( <i>VEV</i> )	0.656 ( <i>VVV</i> )
PCA-EM	0.741	0.910	0.600	0.623	0.391	0.538
kmeans	0.951	0.959	0.866	0.722	0.434	0.502

Table 5: Clustering accuracies on the UCI datasets averaged on 20 trials.

Method	chironomus	wine	iris	zoo	glass	satimage
PCA-kmeans [13]	-	0.702	0.887	0.792	0.472	-
LDA-kmeans [13]	-	0.826	0.980	0.842	0.510	-
Diskmeans [51]	-	-	-	-	-	0.651
DisCluster [51]	-	-	-	-	-	0.642
HMFA [39]	0.987	-	-	-	-	-

Table 6: Clustering accuracies on UCI datasets found in the literature (these results have been obtained with slightly different experimental setups).

other methods for most of the UCI datasets such as chironomus, wine, iris, zoo and satellite image datasets. Concerning the glass dataset, the LDA+kmeans presents apparently the best performances compared to the other methods. However, the results of LDA+kmeans have to be qualified since they were obtained by averaging only 5 trials. Finally, it is interesting from a practical point of view to notice that some DLM models work well in most situations. In particular, the  $\text{DLM}_{[\cdot, \beta]}$  models, in which the variance outside the discriminant subspace is common to all groups, provide very satisfying results for all the datasets considered here.

## 6 Application to mass spectrometry

In this last experimental section, the Fisher-EM procedure is applied to the problem of cancer detection using MALDI mass spectrometry. MALDI mass spectrometry is a non invasive biochemical technique which is useful in searching for disease biomarkers, assessing tumor progression or evaluating the efficiency of drug treatment, to name just a few applications. In particular, a promising field of application is the early detection of the colorectal cancer, which is one of the principal causes of cancer-related mortality, and MALDI imaging could in few years avoid in some cases the colonoscopy method which is invasive and quite expensive.

### 6.1 Data and experimental setup

The MALDI2009 dataset has been provided by Theodore Alexandrov from the Center for Industrial Mathematics (University of Bremen, Germany) and is made of 112 spectra of length 16 331. Among the 112 spectra, 64 are spectra from patients with the colorectal cancer (referred to as cancer hereafter) and 48 are spectra from healthy persons (referred to as control). Each of the 112 spectra is a high-dimensional vector of 16 331 dimensions which covers the mass-to-charge ( $m/z$ ) ratios from 960 to 11 163 Da. For further reading, the dataset is presented in detail and analyzed in a supervised classification framework in [3].

Following the experimental protocol of [3], Fisher-EM was applied on the 6 168 dimensions corresponding to  $m/z$  ratios between 960 and 3 500 Da since there is no discriminative information on the reminder. Figure 8 shows the mean spectra of the cancer and control classes estimated by Fisher-EM on the  $m/z$  interval 900–3500 Da. To be able to compare the clustering results of Fisher-EM, PCA-EM and mixture of PPCA (Mixt-PPCA) have been applied to this subset as well. It has been asked to all methods to cluster the dataset into 2 groups. It is important to remark that this clustering problem is a  $n \ll p$  problem and, among the model-based methods, only these three methods are able to deal with it (see Section 4.5).

### 6.2 Experimental results

Table 7 presents the confusion tables computed from the clustering results of PCA-EM, mixture of PPCA and Fisher-EM. On the one hand, PCA-EM has selected  $d = 4$  principal axes




Figure 8: Estimated mean spectra of the cancer class (up) and of the control class (bottom) on the  $m/z$  interval 900–3500 Da.

PCA-EM			Mixt-PPCA			Fisher-EM		
<i>Cluster</i>			<i>Cluster</i>			<i>Cluster</i>		
<i>Class</i>	Cancer	Control	<i>Class</i>	Cancer	Control	<i>Class</i>	Cancer	Control
Cancer	48	16	Cancer	62	2	Cancer	57	7
Control	1	47	Control	10	38	Control	3	45
<i>Misclassification rate = 0.15</i>			<i>Misclassification rate = 0.11</i>			<i>Misclassification rate = 0.09</i>		

Table 7: Confusion tables for PCA-EM (left), mixture of PPCA (center) and Fisher-EM (right).

with the 90% variance rule before to cluster the data in this subspace and mixture of PPCA has selected  $d = 2$  principal axes for each group. On the other hand, Fisher-EM has estimated the discriminative latent subspace with  $d = K - 1 = 1$  axis to cluster this high-dimensional dataset. It first appears that PCA-EM and mixture of PPCA provide satisfying clustering results on such a complex dataset. However, it is disappointing to see that the PCA-EM make a significant number of false negatives (cancers classified as non-cancers) since the classification risk is not symmetric here. Conversely, mixture of PPCA and Fisher-EM provide a better clustering results both from a global point of view (respectively 89% and 91% of clustering accuracy) and from a medical point of view since Fisher-EM makes significantly less false negatives with an acceptable number of false positives.

More importantly, Fisher-EM provides informations which can be interpreted *a posteriori* to better understand both the data and the phenomenon. Indeed, the values of the estimated loading matrix  $U$ , which is a  $6168 \times 1$  matrix here, expressed the correlation between the discriminative subspace and the original variables. It is therefore possible to identify the original variables with the highest power of discrimination. It is important to highlight that Fisher-EM extracts these informations from the data in a unsupervised framework. Figure 9 shows the correlation between each original variable and the discriminative subspace on an arbitrary scale. The peaks of this curve correspond to the original variables which have a high correlation with the discriminative axis estimated by Fisher-EM.

Figure 10 plots the difference between the mean spectra of the classes cancer and control (cancer - control) and indicates as well, using red triangles, the most discriminative original variables (m/z values). It is not surprising to see that original variables where the cancer and control spectra have a big difference are among the most discriminative. More surprisingly, Fisher-EM selects the original variables with m/z values equal to 2800 and 3050 as discriminative variables whereas the difference between cancer and control spectra is less for these variables than the difference on the variable with m/z value equal to 1350. Such information, which have extracted from the data in a unsupervised framework, may help the pratician to understand the clustering results.

## 7 Conclusion and further works

This work has presented a discriminative latent mixture model which models the data in a latent orthonormal discriminative subspace with an intrinsic dimension lower than the dimension of the original space. A family of 8 parsimonious DLM models has been exhibited by constraining model parameters within and between groups. An estimation algorithm, called the Fisher-EM algorithm, has been also proposed for estimating both the mixture parameters and the latent discriminative subspace. The determination procedure for the discriminative subspace adapts the well-known Fisher criterion to the unsupervised classification context under an orthonormality constraint. Furthermore, when the number of groups is not too






Figure 9: Discrimination power of the original variables: correlation between original variables and the discriminative subspace on an arbitrary scale.




Figure 10: Difference between the mean spectra of the classes cancer and control (cancer - control) and most discriminative variables (indicated by red triangles).

large, the estimated discriminative subspace allows a useful projection of the clustered data. Experiments on simulated and real datasets have shown that Fisher-EM outperforms existing clustering methods. The Fisher-EM algorithm has been also applied to the clustering of mass spectrometry data, which is a real-world and complex application. In this specific context, Fisher-EM has shown its ability to both efficiently cluster high-dimensional mass spectrometry data and give a pertinent interpretation of the results.

Among the possible extensions of this work, it could be first interesting to find a way to visualize in 2D or 3D the clustered data when the estimated discriminative subspace has more than 4 dimensions. Another extension could be to consider a kernel version of Fisher-EM. For this, it would be necessary to replace the Gram matrix introduced in Section 4.5 by a kernel. Finally, it could be also interesting to introduce sparsity in the loading matrix through  $L_1$  penalty.

## A Appendix

In order not to surcharge the notations, the index  $q$  of the current iteration of the Fisher-EM algorithm is not indicated in the following demonstrations. We also define the matrices  $\tilde{W}$  and  $\bar{W}$  such that  $W = \tilde{W} + \bar{W}$ . The matrix  $\tilde{W}$  is defined as a  $p \times p$  matrix containing the  $d$  first vectors of  $W$  completed by zeros such as  $\tilde{W} = [U, 0_{p-d}]$  and  $\bar{W} = W - \tilde{W}$  is defined by  $\bar{W} = [0_d, V]$ .

### A.1 E step

*Proof of Proposition 1.* The conditional expectation  $t_{ik} = E[z_{ik}|y_i, \Theta]$  can be viewed as well as the posterior probability of the observation  $y_i$  given a group  $k$  and, thanks to the Bayes' formula, can be written:

$$t_{ik} = \frac{\pi_k \phi(y_i, \theta_k)}{\sum_{k=1}^K \pi_k \phi(y_i, \theta_k)},$$

where  $\phi$  is the Gaussian density, and  $\pi_k$  and  $\theta_k$  are the parameters of the  $k^{th}$  mixture component estimated in the previous iteration. This posterior probability  $t_{ik}$  can also be formulated from the cost function  $\Gamma_k$  such that:

$$t_{ik} = \frac{1}{\sum_{l=1}^k \exp\left(\frac{1}{2}(\Gamma_k(y_i) - \Gamma_l(y_i))\right)},$$

where  $\Gamma_k(y_i) = -2\log(\pi_k \phi(y_i, \theta_k))$ . According to the assumptions of the DLM models and given that  $W = \tilde{W} + \bar{W}$ ,  $\Gamma_k$  can be reformulated as:

$$\begin{aligned} \Gamma_k(y_i) = & (y_i - m_k)^T \tilde{W} \Delta_k^{-1} \tilde{W}^T (y_i - m_k) + (y_i - m_k)^T \bar{W} \Delta_k^{-1} \bar{W}^T (y_i - m_k) \\ & + \log(|\Delta_k|) - 2\log(\pi_k) + \log(2\pi), \end{aligned}$$

Moreover, since the relations  $\tilde{W}(\tilde{W}^T \tilde{W}) = \tilde{W}$  and  $\bar{W}(\bar{W}^T \bar{W}) = \bar{W}$  hold due to the construction of  $\tilde{W}$  and  $\bar{W}$ , then:

$$\begin{aligned}\Gamma_k(y_i) = & (y_i - m_k)^T \tilde{W} \tilde{W}^T \tilde{W} \Delta_k^{-1} \tilde{W}^T \tilde{W} \tilde{W}^T (y_i - m_k) \\ & + (y_i - m_k)^T \bar{W} \bar{W}^T \bar{W} \Delta_k^{-1} \bar{W}^T \bar{W} \bar{W}^T (y_i - m_k) \\ & + \log(|\Delta_k|) - 2\log(\pi_k) + \log(2\pi),\end{aligned}$$

and:

$$\begin{aligned}\Gamma_k(y_i) = & \left( \tilde{W} \tilde{W}^T (y_i - m_k) \right)^T \tilde{W} \Delta_k^{-1} \tilde{W}^T \left( \tilde{W} \tilde{W}^T (y_i - m_k) \right) \\ & + \frac{1}{\beta_k} \left( \bar{W} \bar{W}^T (y_i - m_k) \right)^T \left( \bar{W} \bar{W}^T (y_i - m_k) \right) \\ & + \log(|\Delta_k|) - 2\log(\pi_k) + \log(2\pi).\end{aligned}$$

Let us now define  $\vartheta_k = \tilde{W} \Delta_k^{-1} \tilde{W}^T$  and  $\|\cdot\|_{\vartheta_k}$ , a norm on the latent space spanned by  $\tilde{W}$ , such that  $\|y\|_{\vartheta_k}^2 = y^T \vartheta_k y$ . With these notations,  $\Gamma_k$  can be rewritten as:

$$\begin{aligned}\Gamma_k(y_i) = & \|\tilde{W} \tilde{W}^T (y_i - m_k)\|_{\vartheta_k}^2 + \frac{1}{\beta_k} \|\bar{W} \bar{W}^T (y_i - m_k)\|^2 \\ & + \log(|\Delta_k|) - 2\log(\pi_k) + \log(2\pi),\end{aligned}$$

and, according to the definition of  $\Delta_k$ :

$$\begin{aligned}\Gamma_k(y_i) = & \|\tilde{W} \tilde{W}^T (y_i - m_k)\|_{\vartheta_k}^2 + \frac{1}{\beta_k} \|\bar{W} \bar{W}^T (y_i - m_k)\|^2 \\ & + \sum_{j=1}^d \log(\alpha_{jk}) + (p-d) \log(\beta_k) - 2\log(\pi_k) + \log(2\pi).\end{aligned}$$

Let us also define the projection operators  $P$  and  $P^\perp$  on the subspaces  $\mathbb{E}$  and  $\mathbb{E}^\perp$  respectively:

- $P(y) = \tilde{W} \tilde{W}^T y$  is the projection of  $y$  on the discriminative space  $\mathbb{E}$ ,
- $P^\perp(y) = \bar{W} \bar{W}^T y$  is the projection of  $y$  on the complementary space  $\mathbb{E}^\perp$ .

Consequently, the cost function  $\Gamma_k$  can be finally reformulated as:

$$\begin{aligned}\Gamma_k(y_i) = & \|P(y_i - m_k)\|_{\vartheta_k}^2 + \frac{1}{\beta_k} \|P^\perp(y_i - m_k)\|^2 \\ & + \sum_{j=1}^d \log(\alpha_{jk}) + (p-d) \log(\beta_k) - 2\log(\pi_k) + \log(2\pi).\end{aligned}$$

Since  $P^\perp(y) = y - P(y)$ , then the distance associated with the complementary subspace can be rewritten as  $\|P^\perp(y_i - m_k)\|^2 = \|(y_i - m_k) - P(y_i - m_k)\|^2$  and this allow to conclude.  $\square$

## A.2 M step

*Proof of Proposition 2.* In the case of the model  $\text{DLM}_{[\alpha_{kj}, \beta_k]}$ , at iteration  $q$ , the conditional expectation of the complete log-likelihood  $Q(y_1, \dots, y_n, \theta | \theta^{(q-1)})$  of the observed data  $\{y_1, \dots, y_n\}$  has the following form:

$$\begin{aligned} Q(\theta) &= \sum_{i=1}^n \sum_{k=1}^K t_{ik} \log(\pi_k \phi(y_i, \theta_k)) \\ &= \sum_{i=1}^n \sum_{k=1}^K t_{ik} \left[ -\frac{1}{2} \log(|S_k|) - \frac{1}{2} (y_i - m_k)^T S_k^{-1} (y_i - m_k) + \log(\pi_k) + \gamma \right], \end{aligned}$$

where  $t_{ik} = E[z_{ik} | \theta^{(q-1)}]$  and  $\gamma = p \log(2\pi)$  is a constant term. According to the definitions of the diagonal matrix  $\Delta_k$  and of the orientation matrix  $W$  for which  $W^{-1} = W^T$ , the inverse covariance matrix  $S_k^{-1}$  of  $Y$  can be written as:

$$S_k^{-1} = (W \Delta_k W^T)^{-1} = W^{-T} \Delta_k^{-1} W^{-1} = W \Delta_k^{-1} W^T.$$

The determinant of  $S_k$  can be also reformulated in the following way:

$$|S_k| = |\Delta_k| = \prod_{j=1}^d \alpha_{kj} \prod_{j=d+1}^p \beta_k = \prod_{j=1}^d \alpha_{jk} \beta_k^{p-d}.$$

Consequently, the  $Q(\theta)$  can be rewritten as:

$$\begin{aligned} Q(\theta) &= -\frac{1}{2} \sum_{k=1}^K n_k \left[ -2 \log(\pi_k) + \sum_{j=1}^d \log(\alpha_{kj}) + (p-d) \log(\beta_k) \right. \\ &\quad \left. + \frac{1}{n_k} \sum_{i=1}^n z_{ik} (y_i - m_k)^T W \Delta_k^{-1} W^T (y_i - m_k) + \gamma \right]. \end{aligned} \tag{A.1}$$

where  $n_k = \sum_{i=1}^n z_{ik}$ . At this point, two remarks can be done on the quantity  $\sum_{i=1}^n z_{ik} (y_i - m_k)^T W \Delta_k^{-1} W^T (y_i - m_k)$ . First, as this quantity is a scalar, it is equal to its trace. Secondly, this quantity can be divided in two parts since  $W = [U, V]$  and  $W = \tilde{W} + \bar{W}$ . Then, the relation  $W \Delta_k^{-1} W^T = \tilde{W} \Delta_k^{-1} \tilde{W}^T + \bar{W} \Delta_k^{-1} \bar{W}^T$  is stated and we can write:

$$\begin{aligned} (y_i - m_k)^T W \Delta_k^{-1} W^T (y_i - m_k) &= \text{tr} \left( (y_i - m_k)^T \tilde{W} \Delta_k^{-1} \tilde{W}^T (y_i - m_k) \right) \\ &\quad + \text{tr} \left( (y_i - m_k)^T \bar{W} \Delta_k^{-1} \bar{W}^T (y_i - m_k) \right). \end{aligned}$$

Moreover, pointing out that  $C_k = \sum_{i=1}^n z_{ik}(y_i - m_k)^T(y_i - m_k)$  is the empirical covariance matrix the  $k$ th group, the previous quantity can be rewritten as:

$$\sum_{i=1}^n z_{ik}(y_i - m_k)^T W \Delta_k^{-1} W^T (y_i - m_k) = \text{tr}(\Delta_k^{-1} \tilde{W}^T C_k \tilde{W}) + \text{tr}(\Delta_k^{-1} \bar{W}^T C_k \bar{W})$$

and finally:

$$\sum_{i=1}^n z_{ik}(y_i - m_k)^T W \Delta_k^{-1} W^T (y_i - m_k) = \sum_{j=1}^d \frac{u_j^T C_k u_j}{\alpha_{kj}} + \sum_{j=d+1}^p \frac{v_j^T C_k v_j}{\beta_k},$$

where  $u_j$ , respectively  $v_j$ , is the  $j$ th column vector of  $U$ , respectively  $V$ . Consequently, replacing this quantity in (A.1) provides the final expression of  $Q(\theta)$ .  $\square$

*Proof of Proposition 3.* The maximization of  $Q(\theta)$  conduces for the DLM models to the following estimates.

**Estimation of  $\pi_k$**  The prior probability  $\pi_k$  of the group  $k$  can be estimated by maximizing  $Q(\theta)$  with respect to the constraint  $\sum_{k=1}^K \pi_k = 1$  which is equivalent to maximize the Lagrange function:

$$L = Q(\theta) + \lambda \left( \sum_{k=1}^K \pi_k - 1 \right),$$

where  $\lambda$  is the Lagrange multiplier. Then, the partial derivative of  $L$  according to  $\pi_k$  is:

$$\frac{\partial L}{\partial \pi_k} = \frac{n_k}{\pi_k} + \lambda.$$

Consequently:

$$\forall k = 1, \dots, K \quad \frac{\partial L}{\partial \pi_k} = 0 \iff \frac{n_k}{\pi_k} + \lambda = 0 \iff n_k + \lambda \pi_k = 0,$$

and:

$$\sum_{k=1}^K (n_k + \lambda \pi_k) = n + \lambda = 0 \implies \lambda = -n.$$

Replacing  $\lambda$  by its value in the partial derivative conduces to an estimation of  $\pi_k$  by:

$$\hat{\pi}_k = \frac{n_k}{n}.$$

**Estimation of  $\mu_k$**  The mean  $\mu_k$  of the  $k$ th group in the latent space can be also estimated by maximizing the expectation of the complete log-likelihood, which can be written in the

following way:

$$Q(\theta) = \sum_{i=1}^n \sum_{k=1}^K t_{ik} \left[ -\frac{1}{2} \log(|S_k|) - \frac{1}{2} (y_i - U\mu_k)^T \Sigma_k^{-1} (y_i - U\mu_k) + \log(\pi_k) \right] + \gamma.$$

Consequently, the partial derivative of  $Q$  according to  $\mu_k$  is:

$$\frac{\partial Q(\theta)}{\mu_k} = \sum_{i=1}^n t_{ik} U^T (y_i - U\mu_k).$$

Setting this quantity to 0 gives:

$$\frac{\partial Q(\theta)}{\mu_k} = 0 \iff \sum_{i=1}^n t_{ik} U^T y_i = \sum_{i=1}^n t_{ik} \mu_k.$$

and conduces to:

$$\hat{\mu}_k = \frac{1}{n_k} \sum_{i=1}^n t_{ik} U^T y_i.$$

**Model** DLM $_{[\alpha_{kj}\beta_k]}$  The partial derivative of  $Q$  according to  $\alpha_{kj}$  is:

$$\frac{\partial Q(\theta)}{\alpha_{kj}} = \frac{1}{\alpha_{kj}} - \frac{u_j^T C_k u_j}{\alpha_{kj}^2},$$

and setting  $\frac{\partial Q(\theta)}{\alpha_{kj}}$  to 0 provides the estimate of  $\alpha_{kj}$ :

$$\hat{\alpha}_{kj} = u_j^T C_k u_j. \quad (\text{A.2})$$

The estimation of  $\beta_k$  is also obtained by maximizing  $Q$  subject to  $\beta_k$ . However, since  $\bar{W} = W - \tilde{W}$ , it is possible to write:

$$\begin{aligned} \frac{n_k}{\beta_k} \sum_{j=d+1}^p w_j^T C_k w_j &= \frac{n_k}{\beta_k} \sum_{j=1}^p w_j^T C_k w_j - \frac{n_k}{\beta_k} \sum_{j=1}^d u_j^T C_k u_j \\ &= \frac{n_k}{\beta_k} \sum_{j=1}^p \text{tr}(w_j w_j^T C_k) - \frac{n_k}{\beta_k} \sum_{j=1}^d u_j^T C_k u_j \\ &= \text{tr} \left( \frac{n_k}{\beta_k} C_k \right) - \frac{n_k}{\beta_k} \sum_{j=1}^d u_j^T C_k u_j \\ &= \frac{n_k}{\beta_k} \left[ \text{tr}(C_k) - \sum_{j=1}^d u_j^T C_k u_j \right]. \end{aligned}$$

Hence:

$$\frac{\partial Q(\theta)}{\beta_k} = 0 \iff \frac{p-d}{\beta_k} - \frac{\text{tr}(C_k)}{\beta_k^2} + \frac{1}{\beta_k^2} \sum_{j=1}^d u_j^T C_k u_j = 0,$$

and it is possible to conclude:

$$\hat{\beta}_k = \frac{\text{tr}(C_k) - \sum_{j=1}^d u_j^T C_k u_j}{p-d}. \quad (\text{A.3})$$

**Model DLM<sub>[\alpha\_{kj}\beta]</sub>** In this case, the variance outside the discriminant subspace is common to all groups and the  $Q(\theta)$  has the following form:

$$\begin{aligned} Q(\theta) &= -\frac{1}{2} \left( \sum_{k=1}^K n_k \left[ -2 \log(\pi_k) + \sum_{j=1}^d \left( \log(\alpha_{kj}) + \frac{u_j^T C_k u_j}{\alpha_{kj}} \right) + \gamma \right] \right. \\ &\quad \left. + \sum_{k=1}^K n_k (p-d) \log(\beta) + \sum_{k=1}^K \frac{n_k}{\beta} \left[ \text{tr}(C_k) - \sum_{j=1}^d u_j^T C_k u_j \right] \right), \\ &= -\frac{1}{2} \left( \sum_{k=1}^K n_k \left[ -2 \log(\pi_k) + \sum_{j=1}^d \left( \log(\alpha_{kj}) + \frac{u_j^T C_k u_j}{\alpha_{kj}} \right) + \gamma \right] \right. \\ &\quad \left. + n(p-d) \log(\beta) + \frac{1}{\beta} \left[ \sum_{k=1}^K \text{tr}(n_k C_k) - \sum_{j=1}^d \sum_{k=1}^K u_j^T n_k C_k u_j \right] \right), \\ &= -\frac{1}{2} \left( \sum_{k=1}^K n_k \left[ -2 \log(\pi_k) + \sum_{j=1}^d \left( \log(\alpha_{kj}) + \frac{u_j^T C_k u_j}{\alpha_{kj}} \right) + \gamma \right] \right. \\ &\quad \left. + n(p-d) \log(\beta) + \frac{1}{\beta} \left[ n \text{tr}(C) - n \sum_{j=1}^d u_j^T C u_j \right] \right), \end{aligned}$$

where  $C$  is the empirical covariance matrix of the whole dataset. Setting to 0 the partial derivative of  $Q(\theta)$  conditionally to  $\beta$  implies:

$$\frac{\partial Q(\theta)}{\beta} = 0 \iff \frac{p-d}{\beta} - \frac{1}{\beta^2} \text{tr}(C) + \frac{1}{\beta^2} \sum_{j=1}^d u_j^T C u_j = 0,$$

and this conduces to:

$$\hat{\beta} = \frac{1}{p-d} \left( \text{tr}(C) - \sum_{j=1}^d u_j^T C u_j \right). \quad (\text{A.4})$$

The estimation of  $\alpha_{kj}$  have been already considered above and is given by Equation (A.2).

**Model DLM<sub>[\alpha\_k \beta\_k]</sub>** For this model, the expectation of the complete log-likelihood  $Q(\theta)$  has the following form:

$$\begin{aligned}
Q(\theta) &= -\frac{1}{2} \sum_{k=1}^K n_k \left[ -2 \log(\pi_k) + \sum_{j=1}^d \left( \log(\alpha_k) + \frac{u_j^T C_k u_j}{\alpha_k} \right) \right. \\
&\quad \left. + (p-d) \log(\beta_k) + \frac{1}{\beta_k} \sum_{j=d+1}^p v_j^T C_k v_j + \gamma \right], \\
Q(\theta) &= -\frac{1}{2} \sum_{k=1}^K n_k \left[ -2 \log(\pi_k) + d \log(\alpha_k) + \frac{1}{\alpha_k} \sum_{j=1}^d u_j^T C_k u_j \right. \\
&\quad \left. + (p-d) \log(\beta_k) + \frac{1}{\beta_k} \sum_{j=d+1}^p v_j^T C_k v_j + \gamma \right].
\end{aligned}$$

The partial derivative of  $Q(\theta)$  according to  $\alpha_k$  is:

$$\frac{\partial Q(\theta)}{\alpha_k} = \frac{d}{\alpha_k} - \frac{1}{\alpha_k^2} \sum_{j=1}^d u_j^T C_k u_j,$$

and, setting this quantity to 0, provides:

$$\hat{\alpha}_k = \frac{1}{d} \sum_{j=1}^d u_j^T C_k u_j. \tag{A.5}$$

On the other hand, the estimation of  $\beta_k$  is the same as in Equation (A.3).

**Model DLM<sub>[\alpha\_k \beta]</sub>** The estimations of  $\alpha_k$  and  $\beta$  are respectively provided by Equations (A.5) and (A.4).



**Model DLM** <sub>$[\alpha_j \beta_k]$</sub>  In this case,  $Q(\theta)$  has the following form:

$$\begin{aligned}
Q(\theta) &= -\frac{1}{2} \sum_{k=1}^K n_k \left( -2 \log(\pi_k) + \sum_{j=1}^d \left( \log(\alpha_j) + \frac{u_j^T C_k u_j}{\alpha_j} \right) \right. \\
&\quad \left. + (p-d) \log(\beta_k) + \frac{1}{\beta_k} \sum_{j=d+1}^p v_j^T C_k v_j + \gamma \right), \\
Q(\theta) &= -\frac{1}{2} \left( \sum_{k=1}^K n_k \left[ -2 \log(\pi_k) \right] + \sum_{k=1}^K n_k \sum_{j=1}^d \log(\alpha_j) + \sum_{j=1}^d \sum_{k=1}^K \frac{u_j^T (n_k C_k) u_j}{\alpha_j} \right. \\
&\quad \left. + \sum_{k=1}^K n_k \left[ (p-d) \log(\beta_k) + \frac{1}{\beta_k} \sum_{j=d+1}^p v_j^T C_k v_j + \gamma \right] \right), \\
Q(\theta) &= -\frac{1}{2} \left( \sum_{k=1}^K n_k \left[ -2 \log(\pi_k) \right] + n \sum_{j=1}^d \log(\alpha_j) + n \sum_{j=1}^d \frac{u_j^T C u_j}{\alpha_j} \right. \\
&\quad \left. + \sum_{k=1}^K n_k \left[ (p-d) \log(\beta_k) + \frac{1}{\beta_k} \sum_{j=d+1}^p v_j^T C_k v_j + \gamma \right] \right).
\end{aligned}$$

The partial derivative of  $Q(\theta)$  according to  $\alpha_j$  is:

$$\frac{\partial Q(\theta)}{\alpha_j} = \frac{1}{\alpha_j} - \frac{1}{\alpha_j^2} u_j^T C u_j,$$

and  $\frac{\partial Q(\theta)}{\alpha_j} = 0$  implies:

$$\hat{\alpha}_j = u_j^T C u_j. \tag{A.6}$$

**Model DLM** <sub>$[\alpha_j \beta]$</sub>  The estimations of  $\alpha_j$  and  $\beta$  are respectively provided by Equations (A.6) and (A.4).

**Model DLM<sub>[\alpha\beta\_k]</sub>** In this case,  $Q(\theta)$  has the following form:

$$\begin{aligned}
Q(\theta) &= -\frac{1}{2} \sum_{k=1}^K n_k \left( -2 \log(\pi_k) + d \log(\alpha) + \frac{1}{\alpha} \sum_{j=1}^d u_j^T C_k u_j + \right. \\
&\quad \left. + (p-d) \log(\beta_k) + \frac{1}{\beta_k} \sum_{j=d+1}^p v_j^T C_k v_j + \gamma \right), \\
Q(\theta) &= -\frac{1}{2} \left( \sum_{k=1}^K n_k [-2 \log(\pi_k)] + \sum_{k=1}^K n_k d \log(\alpha) + \frac{1}{\alpha} \sum_{j=1}^d \sum_{k=1}^K n_k u_j^T C_k u_j \right. \\
&\quad \left. + \sum_{k=1}^K n_k \left[ (p-d) \log(\beta_k) + \frac{1}{\beta_k} \sum_{j=d+1}^p v_j^T C_k v_j + \gamma \right] \right), \\
Q(\theta) &= -\frac{1}{2} \left( \sum_{k=1}^K n_k [-2 \log(\pi_k)] + n \log(\alpha) + \frac{n}{\alpha} \sum_{j=1}^d u_j^T C u_j \right. \\
&\quad \left. + \sum_{k=1}^K n_k \left[ (p-d) \log(\beta_k) + \frac{1}{\beta_k} \sum_{j=d+1}^p v_j^T C_k v_j + \gamma \right] \right),
\end{aligned}$$

The partial derivative of  $Q(\theta)$  according to  $\alpha$  is:

$$\frac{\partial Q(\theta)}{\partial \alpha} = \frac{d}{\alpha} - \frac{1}{\alpha^2} \sum_{j=1}^d u_j^T C u_j,$$

and, setting this quantity to 0, we end up with:

$$\hat{\alpha} = \frac{1}{d} \sum_{j=1}^d u_j^T C u_j. \quad (\text{A.7})$$

The estimation of  $\beta_k$  is the same as in Equation (A.3).

**Model DLM<sub>[\alpha\beta]</sub>** The estimations of  $\alpha$  and  $\beta$  have been already computed and are provided by Equations (A.7) and (A.4).  $\square$

## References

- [1] R. Agrawal, J. Gehrke, D. Gunopulos, and P. Raghavan. Automatic subspace clustering of high-dimensional data for data mining application. In *ACM SIGMOD International Conference on Management of Data*, pages 94–105, 1998.
- [2] H. Akaike. A new look at the statistical model identification. *IEEE Transactions on Automatic Control*, 19(6):716–723, 1974.

- [3] T. Alexandrov, J. Decker, B. Mertens, A.M. Deelder, R.A. Tollenaar, P. Maass, and H. Thiele. Biomarker discovery in MALDI-TOF serum protein profiles using discrete wavelet transformation. *Bioinformatics*, 25(5):643–649, 2009.
- [4] E. Anderson. The irises of the gaspé peninsula. *Bulletin of the American Iris Society*, 59:2–5, 1935.
- [5] J. Baek, G. McLachlan, and L. Flack. Mixtures of Factor Analyzers with Common Factor Loadings: Applications to the Clustering and Visualisation of High-Dimensional Data. *IEEE Transactions on Pattern Analysis and Machine Intelligence*, pages 1–13, 2009.
- [6] R. Bellman. *Dynamic Programming*. Princeton University Press, 1957.
- [7] C. Biernacki, G. Celeux, and G. Govaert. Assessing a mixture model for clustering with the integrated completed likelihood. *IEEE Transactions on Pattern Analysis and Machine Intelligence*, 22(7):719–725, 2001.
- [8] C. Biernacki, G. Celeux, and G. Govaert. Choosing starting values for the EM algorithm for getting the highest likelihood in multivariate Gaussian mixture models. *Computational Statistics and Data Analysis*, 41:561–575, 2003.
- [9] C. Bouveyron, S. Girard, and C. Schmid. High dimensional discriminant analysis. *Communication in Statistics: Theory and Methods*, 52(1):502–519, 2007.
- [10] G. Celeux and J. Diebolt. The sem algorithm: a probabilistic teacher algorithm from the em algorithm for the mixture problem. *Computational Statistics Quarterly*, 2(1):73–92, 1985.
- [11] G. Celeux and G. Govaert. A classification em algorithm for clustering and two stochastic versions. *Computational Statistics and Data Analysis*, 14:315–332, 1992.
- [12] A. Dempster, N. Laird, and D. Robin. Maximum likelihood from incomplete data via the em algorithm. *Journal of the Royal Statistical Society*, 39(1):1–38, 1977.
- [13] C. Ding and T. Li. Adaptive dimension reduction using discriminant analysis and k-means clustering. *ICML*, 2007.
- [14] R. Duda, P. Hart, and D. Stork. *Pattern classification*. John Wiley & Sons, 2000.
- [15] R.A. Fisher. The use of multiple measurements in taxonomic problems. *Annals of Eugenics*, 7:179–188, 1936.
- [16] D.H. Foley and J.W. Sammon. An optimal set of discriminant vectors. *IEEE Transactions on Computers*, 24:281–289, 1975.

- [17] C. Fraley and A. Raftery. Model-based clustering, discriminant analysis, and density estimation. *Journal of the American Statistical Association*, 97(458), 2002.
- [18] J.H. Friedman. Regularized discriminant analysis. *The journal of the American statistical association*, 84:165–175, 1989.
- [19] K. Fukunaga. *Introduction to Statistical Pattern Recognition*. Academic. Press, San Diego, 1990.
- [20] S. Girard. A non-linear pca based on manifold approximation. *Computational Statistics*, 15(2):145–167, 2000.
- [21] Y-F. Guo, S-J. Li, J-Y. Yang, T-T. Shu, and L-D. Wu. A generalized foley-sammon transform based on generalized fisher discriminant criterion and its application to face recognition. *Pattern Recognition letters*, 24:147–158, 2003.
- [22] Y. Hamamoto, Y. Matsuura, T. Kanaoka, and S. Tomita. A note on the orthonormal discriminant vector method for feature extraction. *Pattern Recognition*, 24(7):681–684, 1991.
- [23] T. Hastie, A. Buja, and R. Tibshirani. Penalized discriminant analysis. *Annals of Statistics*, 23:73–102, 1995.
- [24] T. Hastie and W. Stuetzle. Principal curves. *Journal of the American Statistical Association*, 84:502–516, 1989.
- [25] T. Hastie, R. Tibshirani, and J. Friedman. *The elements of statistical learning*. Springer, New York, second edition, 2009.
- [26] P. Howland and H. Park. Generalizing discriminant analysis using the generalized singular decomposition. *IEEE transactions on pattern analysis and machine learning*, 2004.
- [27] A. Jain, M. Marty, and P. Flynn. Data Clustering: a review. *ACM Computing Surveys*, 31(3):264–323, 1999.
- [28] Z. Jin, J.Y. Yang, Z.S. Hu, and Z. Lou. Face recognition based on the uncorrelated optimal discriminant vectors. *Pattern Recognition*, 10(34):2041–2047, 2001.
- [29] I. Jolliffe. *Principal Component Analysis*. Springer-Verlag, New York, 1986.
- [30] N. Kambhatla and T. Leen. Dimension reduction by local principal component analysis. *Neural Computation*, 9:1493–1516, 1997.
- [31] T. Kohonen. *Self Organizing Maps*. Springer, Berlin, heidleberg edition, 1995.
- [32] F. De la Torre Frade and T. Kanade. Discriminative cluster analysis. *ICML*, pages 241–248, 2006.

- [33] K. Liu, Y-Q. Cheng, and J-Y. Yang. A generalized optimal set of discriminant vectors. *Pattern Recognition*, 25(7):731–739, 1992.
- [34] C. Maugis, G. Celeux, and M.-L. Martin-Magniette. Variable selection for Clustering with Gaussian Mixture Models. *Biometrics*, 65(3):701–709, 2009.
- [35] G. McLachlan and T. Krishnan. *The EM algorithm and extensions*. Wiley Interscience, New York, 1997.
- [36] G. McLachlan and D. Peel. *Finite Mixture Models*. Wiley Interscience, New York, 2000.
- [37] G. McLachlan, D. Peel, and R. Bean. Modelling high-dimensional data by mixtures of factor analyzers. *Computational Statistics and Data Analysis*, (41):379.
- [38] P. McNicholas and B. Murphy. Parsimonious Gaussian mixture models. *Statistics and Computing*, 18(3):285–296, 2008.
- [39] A. Montanari and C. Viroli. Heteroscedastic Factor Mixture Analysis. *Statistical Modeling: An International journal (forthcoming)*, (to appear), 2010.
- [40] L. Parsons, E. Haque, and H. Liu. Subspace clustering for high dimensional data: a review. *SIGKDD Explor. Newsl.*, 6(1):69–76, 1998.
- [41] A. Raftery and N. Dean. Variable selection for model-based clustering. *Journal of the American Statistical Association*, 101(473):168–178, 2006.
- [42] D. Rubin and D. Thayer. Em algorithms for ml factor analysis. *Psychometrika*, 47(1):69–76, 1982.
- [43] B. Schölkopf, A. Smola, and K. Müller. Non linear component analysis as a kernel eigenvalue problem. *Neural Computation*, 10:1299–1319, 1998.
- [44] G. Schwarz. Estimating the dimension of a model. *The Annals of Statistics*, 6:461–464, 1978.
- [45] D. Scott and J. Thompson. Probability density estimation in higher dimensions. In *Fifteenth Symposium in the Interface*, pages 173–179, 1983.
- [46] E. Tipping and C. Bishop. Mixtures of probabilistic principal component analysers. *Neural Computation*, 11(2):443–482, 1999.
- [47] N. Trendafilov and I. T. Jolliffe. DALASS: Variable selection in discriminant analysis via the LASSO. *Computational Statistics and Data Analysis*, 51:3718–3736, 2007.
- [48] M. Verleysen and D. François. The curse of dimensionality in data mining and time series prediction. *IWANN*, 2005.

- [49] M. Xu and P. Fränti. Iterative k-means algorithm based on fisher discriminant. 1, 1997.
- [50] J. Ye. Characterization of a family of algorithms for generalized discriminant analysis on undersampled problems. *Journal of Machine Learning Research*, 6:483–502, 2005.
- [51] J. Ye, Z. Zhao, and M. Wu. Discriminative k-means for clustering. *Advances in Neural Information Processing Systems 20*, pages 1649–1656, 2007.
- [52] Z. Zhang, G. Dai, and M.I. Jordan. A flexible and efficient algorithm for regularized fisher discriminant analysis, 2009.